arXiv:math/0409025v1 [math.OA] 2 Sep 2004

CUMULANTS IN NONCOMMUTATIVE PROBABILITY THEORY IV. NONCROSSING CUMULANTS: DE FINETTI'S THEOREM, L^p-INEQUALITIES AND BRILLINGER'S FORMULA

FRANZ LEHNER

Abstract. In this paper we collect a few results about exchangeability systems in which crossing cumulants vanish, which we call noncrossing exchangeability systems.

De Finetti's theorem states that any exchangeable sequence of random variables is conditionally i.i.d. with respect to some σ-algebra. In this paper we prove a "free" noncommutative analog of this theorem, namely we show that any noncrossing exchangeability system with a faithful state which satisfies a so called weak singleton condition, can be embedded into a free product with amalgamation over a certain subalgebra. Vanishing of crossing cumulants can be verified by checking a certain weak freeness condition. The proof follows the classical proof of De Finetti's theorem, the main technical tool being a noncommutative L^p-inequality for i.i.d. sums of centered noncommutative random variables in noncrossing exchangeability systems.

In the second part of the paper we establish a free analogue of Brillinger's formula, which expresses free cumulants in terms of amalgamated free cumulants.

Contents

1. Introduction 2
2. Preliminaries 3
2.1. Exchangeability Systems and Cumulants 3
2.2. Noncrossing partitions and freeness 4
2.3. Multiplicative functions and convolution on the lattice of noncrossing partitions 5
2.4. The weak singleton condition 7
2.5. Conditional Expectations 8
2.6. Statement of main result 10
3. A de Finetti Lemma 10
4. A noncommutative L^p-inequality in the case of noncrossing cumulants 16
5. Weak freeness 20
6. Appendix: Free L^p Inequalities 24
7. A Free Analog of Brillinger's Formula 25
7.1. Brillinger's formula for classical cumulants 25
7.2. Free cumulants 26
7.3. Conditioned free cumulants 26
7.4. An application 27

Date: February 1, 2008.

1991 *Mathematics Subject Classification.* Primary 46L53, Secondary 05A18.

Key words and phrases. Cumulants, partition lattice, Möbius inversion, free probability, noncrossing partitions, noncommutative probability.

Supported by the Austrian Science Fund (FWF) Project №R2-MAT.

Acknowledgements 30
References 30

1. INTRODUCTION

De Finetti's theorem [Kin78, CT78] states that for any exchangeable sequence of random variables there exists a σ-algebra conditionally on which the sequence is i.i.d. There are various noncommutative versions of this theorem [Stø69, HM76, Hud81, Pet90, AL93], all of which involve tensor product constructions or other commutativity conditions. Indeed there is no hope to obtain a general De Finetti's theorem without imposing additional conditions.

In this paper we consider conditions under which a kind of "most noncommutative" version of De Finetti's theorem holds, namely a characterization of exchangeability systems which can be written as amalgamated free products.

The only prerequisite for this paper is part I of the series [Leh04], where exchangeability systems are introduced and many examples are discussed. In section I.4.5 of that paper we discussed the amalgamated free product as an operator valued exchangeability system with a conditional expectation ψ. Composing this conditional expectation with a state on the amalgamated subalgebra gives rise to a scalar valued exchangeability system.

The question now is, given an exchangeability system $\mathcal{E}$, when is it of this form?

An obvious necessary condition is that crossing cumulants must vanish, because this is the case for the operator-valued amalgamated free cumulants and the $\mathcal{E}$-cumulants are simply the expectations of the latter, see section I.3.6. Another necessary condition is a certain *weak singleton condition*. The singleton condition introduced in [BS96] is too strong, because together with the vanishing of crossing cumulants it actually implies freeness. The weak singleton condition however, to be defined below, turns out to be the right one and is automatically satisfied if the state is tracial.

The construction of the conditional expectation essentially follows the classical proof, namely by adjoining the algebra $\mathcal{B}$ of permutation invariant random variables to the initial algebra and extending to it the expectation functional. Moreover there it is possible to construct a conditional expectation ψ_∞ onto $\mathcal{B}$ as the limit of symmetrizing maps ψ_N. There are certain technical issues regarding the faithfulness of the extension in the non-tracial case. These are solved by a certain L^p-inequality which is of some independent interest.

Finally we prove a free analog of Brillinger's formula expressing (scalar) free cumulants in terms of amalgamated free cumulants.

The paper is organized as follows.

In Section 2 we collect a few definitions and lemmas needed for the statement and the proof of the main result.

In Section 3 we adapt a proof from [AL93] to the noncrossing situation. It shows that the conditional expectations ψ_N evaluated at words of the form $X_1^{(h(1))}X_2^{(h(2))}\cdots X_n^{(h(n))}$ asymptotically factor according to the connected components of the kernel partition $\ker h$.

In Section 4 we prove a strong law of large numbers in noncrossing exchangeability systems under the assumption of the weak singleton condition.

In Section 5 we discuss a certain weak freeness condition and show that together with the weak singleton condition it implies that crossing cumulants vanish and thus weak freeness is the same as freeness with amalgamation.

In the final section 7, which can be read independently from the rest of the paper, we prove a free analog of Brillinger's formula [Bri69]. The latter expresses classical cumulants as a sum over set partitions of "cumulants of cumulants". In this section, which is independent of the rest of the paper, we present a certain free analog of this result. As expected, the sum now runs over noncrossing partitions.

2. Preliminaries

In this section we collect the necessary definitions and auxiliary results needed later on. For details we refer to part I [Leh04].

2.1. **Exchangeability Systems and Cumulants.** We recall first that an *exchangeability system* $\mathcal{E}$ for a noncommutative probability space $(\mathcal{A},\varphi)$ consists of another noncommutative probability space $(\mathcal{U},\tilde{\varphi})$ and an infinite family $\mathcal{J}=(\iota_k)_{k\in\mathbf{N}}$ of state-preserving embeddings $\iota_k:\mathcal{A}\to\mathcal{A}_k\subseteq\mathcal{U}$, which we conveniently denote by $X\mapsto X^{(k)}$, such that the algebras $\mathcal{A}_j$ are *interchangeable* with respect to $\tilde{\varphi}$: for any family $X_1,X_2,\dots,X_n\in\mathcal{A}$, and for any choice of indices $h(1),\dots,h(n)$ the expectation is invariant under any permutation $\sigma\in\mathfrak{S}_\infty$ in the sense that

$$\tilde{\varphi}(X_1^{(h(1))}X_2^{(h(2))}\cdots X_n^{(h(n))})=\tilde{\varphi}(X_1^{(\sigma(h(1)))}X_2^{(\sigma(h(2)))}\cdots X_n^{(\sigma(h(n)))}). \tag{2.1}$$

Throughout this paper we will assume that the algebra $\mathcal{U}$ is generated by the algebras $\mathcal{A}_k$ and that the action of $\mathfrak{S}_\infty$ extends to all of $\mathcal{U}$.

Denote by Π_n the lattice of partitions (or equivalence relations) of the set $[n]=\{1,2,\dots,n\}$. The refinement order $\pi\leq\sigma$ means as usual that the partition π is finer than the partition σ.

Then permutation invariance means that the value of the expectation (2.1) only depends on the so-called *kernel* $\pi=\ker h\in\Pi_n$ of the map h which is the partition defined by

$$i\sim_\pi j \qquad\Longleftrightarrow\qquad h(i)=h(j)$$

and we denote the common value as

$$\varphi_\pi(X_1,X_2,\dots,X_n)=\tilde{\varphi}(X_1^{(\pi(1))}X_2^{(\pi(2))}\cdots X_n^{(\pi(n))}). \tag{2.2}$$

Here we consider a partition $\pi\in\Pi_n$ as a function $\pi:[n]\to\mathbf{N}$, mapping each element to the number of the block containing it. This is a canonical example of an index function h with $\ker h=\pi$ and because of condition (2.1) the actual numbering of the blocks does not matter.

Subalgebras $\mathcal{B},\mathcal{C}\subseteq\mathcal{A}$ are called *$\mathcal{E}$-exchangeable* or, more suggestively, *$\mathcal{E}$-independent* if for any choice of random variables $X_1,X_2,\dots,X_n\in\mathcal{B}\cup\mathcal{C}$ and subsets $I,J\subseteq\{1,\dots,n\}$ such that $I\cap J=\emptyset$, $I\cup J=\{1,\dots,n\}$, $X_i\in\mathcal{B}$ for $i\in I$ and $X_i\in\mathcal{C}$ for $i\in J$, we have the identity

$$\varphi_\pi(X_1,X_2,\dots,X_n)=\varphi_{\pi'}(X_1,X_2,\dots,X_n)$$

whenever π, $\pi'\in\Pi_n$ are partitions with $\pi|_I=\pi'|_I$ and $\pi|_J=\pi'|_J$. Two families of random variables (X_i) and (Y_j) are called $\mathcal{E}$-exchangeable if the algebras they generate have this property.

We say that two random variables X and $Y\in\mathcal{A}$ *have the same distribution given $\mathcal{E}$*, if for any word $W=W_1W_2\cdots W_n$ with $W_i\in\{X^{(1)}\}\cup\bigcup_{i\geq2}\mathcal{A}_i$ the expectation $\tilde{\varphi}(W)$

does not change if we replace each occurrence of $X^{(1)}$ by $Y^{(1)}$. We call X and Y *$\mathcal{E}$-i.i.d.* if in addition they are $\mathcal{E}$-independent. Similarly a sequence $(X_i)_{i\in\mathbf{N}} \subseteq \mathcal{A}$ of $\mathcal{E}$-independent random variables is called *$\mathcal{E}$-i.i.d.* if for any word $W = W_1W_2\cdots W_n$ with $W_i \in \{X_i^{(1)} : i \in \mathbf{N}\} \cup \bigcup_{i\geq 2}\mathcal{A}_i$ the expectation $\tilde{\varphi}(W)$ does not change if we apply a permutation $\sigma \in \mathfrak{S}_\infty$ to the indices of X_i, i.e., if we replace each occurrence of X_i by $X_{\sigma(i)}$. Then it is possible to define cumulant functionals, indexed by set partitions $\pi \in \Pi_n$, via

$$K^{\mathcal{E}}_\pi(X_1, X_2, \dots, X_n) = \sum_{\sigma\leq\pi} \varphi_\sigma(X_1, X_2, \dots, X_n)\,\mu(\sigma, \pi)$$

where $\mu(\sigma, \pi)$ is the Möbius function of the lattice of set partitions, cf. part I. The use of the probabilistic termini "independence" and "cumulants" is justified by the following proposition which establishes the analogy to classical probability.

Proposition 2.1 ([Leh04]). *Two subalgebras $\mathcal{B}, \mathcal{C} \subseteq \mathcal{A}$ are $\mathcal{E}$-independent if and only if mixed cumulants vanish, that is, whenever $X_i \in \mathcal{B} \cup \mathcal{C}$ are some noncommutative random variables and $\pi \in \Pi_n$ is an arbitrary partition such that there is a block of π which contains indices i and j such that $X_i \in \mathcal{B}$ and $X_j \in \mathcal{C}$, then $K^{\mathcal{E}}_\pi(X_1, X_2, \dots, X_n)$ vanishes.*

With this abstract formalism we can now transfer many combinatorial proofs from classical probability to the general situation. One of the most useful results is the product formula of Leonov and Shiryaev.

Proposition 2.2 ([Leh04, Prop. 3.3]). *Let $(X_{i,j})_{i\in\{1,\dots,m\}, j\in\{1,\dots,n_i\}} \subseteq \mathcal{A}$ be a family of noncommutative random variables, in total $n = n_1 + n_2 + \cdots + n_m$ variables. Then every partition $\pi \in \Pi_m$ induces a partition $\tilde{\pi}$ on $\{1, \dots, n\} \simeq \{(i, j) : i \in [m], j \in [n_i]\}$ with blocks $\tilde{B} = \{(i, j) : i \in B, j \in [n_i]\}$, that is, each block $B \in \pi$ is replaced by the union of the intervals $(\{n_{i-1} + 1, n_{i-1} + 2, \dots, n_i\})_{i\in B}$. Then we have*

$$K^{\mathcal{E}}_\pi(\prod_{j_1} X_{1,j_1}, \prod_{j_2} X_{2,j_2}, \dots, \prod_{j_m} X_{m,j_m}) = \sum_{\substack{\sigma\in\Pi_n\\ \sigma\vee\tilde{\hat{0}}_m=\tilde{\pi}}} K^{\mathcal{E}}_\sigma(X_{1,1}, X_{1,2}, \dots, X_{m,n_m})$$

Remark 2.3. In the sequel we will sometimes not distinguish between i.i.d. sequences in $\mathcal{A}$ and sequences of the form $X^{(i)}$. The latter do not belong to $\mathcal{A}$ strictly speaking, but we can replace $\mathcal{A}$ by the algebra $\tilde{\mathcal{A}}$ generated by $(\mathcal{A}_i)_{i\in I}$, where $I \subseteq \mathbf{N}$ is an infinite subset, and construct an exchangeability system for $\tilde{\mathcal{A}}$ by considering $\mathbf{N}$ as a disjoint union of infinitely many copies of I.

The constructions above can be done in the more general situation of an *operator valued noncommutative probability space*, which is a pair $(\mathcal{A}, \psi)$ consisting of a unital algebra $\mathcal{A}$ and a *conditional expectation* ψ onto some unital subalgebra $\mathcal{B} \subseteq \mathcal{A}$. A conditional expectation is a unital positive map $\psi : \mathcal{A} \to \mathcal{B}$, with the property that $\psi(BXB') = B\psi(X)B'$ whenever $B, B' \in \mathcal{B}$ and $X \in \mathcal{A}$.

2.2. **Noncrossing partitions and freeness.** The lattice of noncrossing partitions, denoted by NC_n, will play a prominent rôle in this paper. We recall that a partition $\pi \in \Pi_n$ is *noncrossing*, if there is no quadruple of indices $i < j < k < l$, such that $i \sim_\pi k$ and $j \sim_\pi l$ and $i \not\sim_\pi j$. Equivalently, noncrossing partitions can also be characterized by the fact that recursively there is always one block which is an interval and after removing this block the remaining partition is still noncrossing.

Definition 2.4. We say that *crossing cumulants vanish* in a given exchangeability system $\mathcal{E}$ if for any n and for any choice of random variables X_1, X_2,..., X_n the cumulant

$$K_\pi^{\mathcal{E}}(X_1, X_2, \dots, X_n) = 0$$

whenever π has a crossing, i.e., $\pi \notin NC_n$. We call such an exchangeability system a *noncrossing exchangeability system* .

A prominent example of a noncrossing exchangeability system is the free exchangeability system, where $(\mathcal{U}, \tilde{\varphi})$ is the reduced free product of an infinite family of copies of a given noncommutative probability space $(\mathcal{A}, \varphi)$, see Section I.4.4. We recall that the free exchangeability system is characterized by the property that

$$\varphi(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = 0$$

whenever $\varphi(X_j) = 0$ and $h(j) \neq h(j+1)$ for every $1 \leq j \leq n-1$.
Another situation where crossing cumulants vanish is *freeness with amalgamation* [Voi95, Spe98], which is a noncommutative analog of conditional independence. Let $(\mathcal{A}, \psi)$ be a $\mathcal{B}$-valued noncommutative probability space. Then the amalgamated free product $\mathcal{U} = \bigstar_{\mathcal{B}} \mathcal{A}_i$ of infinitely many copies of $\mathcal{A}$ with amalgamation over $\mathcal{B}$ is a $\mathcal{B}$-valued noncrossing exchangeability system for $(\mathcal{A}, \psi)$. The amalgamated free exchangeability system is characterized by the property that

$$\psi(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = 0$$

whenever $\psi(X_j) = 0$ and $h(j) \neq h(j+1)$ for every $1 \leq j \leq n-1$. As in the case of scalar freeness, the $\mathcal{B}$-valued cumulants $K_\pi^{\mathcal{E},\psi}(X_1, \dots, X_n)$ vanish for any partition $\pi \notin NC_n$.
Now choose any state on $\mathcal{B}$, then

$$\mathcal{E}^\varphi = (\mathcal{U} = \bigstar_{\mathcal{B}} \mathcal{A}_i, \varphi \circ \psi, \mathcal{J}) \tag{2.3}$$

is a scalar-valued exchangeability system for the noncommutative probability space $(\mathcal{A}, \varphi \circ \psi)$ whose cumulants are

$$K_\pi^{\mathcal{E},\varphi\circ\psi}(X_1, \dots, X_n) = \varphi(K_\pi^{\mathcal{E},\psi}(X_1, \dots, X_n));$$

again cumulants vanish for any partition π with crossings. Other examples of noncrossing exchangeability systems can be constructed by taking conditionally free products [BLS96]. We will see later that these cannot be realized amalgamated free products, cf. Remark 2.16

2.3. **Multiplicative functions and convolution on the lattice of noncrossing partitions.** We refer to [Sta86] or section I.1.3 for the definition of the incidence algebra of a poset. In the case of noncrossing partitions there is also a reduced incidence algebra of multiplicative functions. Let NCI_n be the set of intervals in NC_n and $NCI = \bigcup_n NCI_n$. Then every interval $[\pi, \sigma]$ has a canonical decomposition

$$[\pi, \sigma] \simeq [\hat{0}_1, \hat{1}_1]^{k_1} \times [\hat{0}_2, \hat{1}_2]^{k_2} \times \cdots \times [\hat{0}_1, \hat{1}_n]^{k_n} \times \cdots \tag{2.4}$$

where $(k_n)_{n=1}^\infty$ is a sequence of integers with finitely many nozero entries and (2.4) is a lattice isomorphism [Spe94]. For example it is easy to see that for $\pi \in NC_n$ we have

$$[\hat{0}_n, \pi] \simeq \prod_{p=1}^n [\hat{0}_p, \hat{1}_p]^{k_p} \tag{2.5}$$

where k_p is the number of blocks of π of size p.

A function $f : NCI \to \mathbf{C}$ is called *multiplicative* if for any interval $[\pi, \sigma]$ it satisfies

$$f([\pi,\sigma]) = f([\hat{0}_1, \hat{1}_1])^{k_1} f([\hat{0}_2, \hat{1}_2])^{k_2} \cdots f([\hat{0}_1, \hat{1}_n])^{k_n}$$

where $[\pi, \sigma]$ has the decomposition (2.4). Such a function is determined by its *characteristic sequence* $f_n = f([\hat{0}_n, \hat{1}_n])$ and it is easy to see that the convolution of two multiplicative functions is again multiplicative, i.e., the multiplicative functions constitute an algebra, the so-called *reduced incidence algebra*. For a noncrossing partition π with decomposition as in (2.5) we will denote

$$f_\pi := f([\hat{0}_n, \pi]) = \prod_{p=1}^{n} f_p^{k_p}.$$

Then the convolution $f \boxtimes g$ of two multiplicative functions can be calculated with the aid of the *Kreweras complementation map* [Kre72]. This is a lattice anti-automorphism of NC_n described as follows. Paint n points on a circle and label them clockwise with numbers 1, 2, …, n. A noncrossing partition $\pi \in NC_n$ can be visualized by drawing inside the circle for each block of π the convex polygon whose vertices are the elements of the block. Now put another n points with labels $\bar{1}$, $\bar{2}$, …, $\bar{n}$ on the circle, placing the point with label $\bar{k}$ between the points with label k and $k+1$ and connect the new points with each other by drawing as many lines as possible without intersecting the polygons drawn before. This leads to a noncrossing partition of the set $\{\bar{1}, \bar{2}, \dots, \bar{n}\}$ which is called the *Kreweras complement* of π and is denoted $K(\pi)$. It is easy to see that K is an order anti-automorphism, $K(\hat{0}_n) = \hat{1}_n$ and $K(\hat{1}_n) = \hat{0}_n$. It follows that $[\pi, \hat{1}_n] \simeq [\hat{0}_n, K(\pi)]$ and therefore the convolution of multiplicative functions can be written

$$(f \boxtimes g)_n = f \boxtimes g([\hat{0}_n, \hat{1}_n]) = \sum_{\pi \in NC_n} f_\pi g_{K(\pi)}.$$

As a consequence the reduced incidence algebra is commutative and there is a "Fourier Transform" [NS97]: Let $(f_n)_{n=1}^\infty$, $(g_n)_{n=1}^\infty$ be the characteristic sequences of two multiplicative functions f and g with $f_1 = g_1 = 1$. The formal power series

$$\varphi_f(z) = \sum_{n=1}^{\infty} f_n z^n$$

is called the *characteristic series* of f. Let $\varphi_f^{\langle -1 \rangle}(z)$ be the compositional inverse of φ_f and

$$\mathcal{F}_f(z) = \frac{1}{z} \varphi_f^{\langle -1 \rangle}(z),$$

then

$$\mathcal{F}_{f \boxtimes g}(z) = \mathcal{F}_f(z)\, \mathcal{F}_g(z) \tag{2.6}$$

Two prominent multiplicative functions are the *Zeta function* $\zeta(\pi, \sigma) \equiv 1$ with characteristic series

$$\varphi_\zeta(z) = \frac{z}{1-z}$$

and its inverse, the *Möbius function* μ whose characteristic sequence is given by the signed Catalan numbers $\mu_n = (-1)^{n-1} C_{n-1} = \frac{(-1)^{n-1}}{n} \binom{2n-2}{n-1}$ and the characteristic series is

$$\varphi_\mu(z) = \frac{\sqrt{1+4z} - 1}{2}.$$

These functions satisfy $\zeta \boxtimes \mu = \delta$, where δ is the unit element of the reduced incidence algebra and has characteristic series $\varphi_\delta(z) = z$. Moreover, functions f and g satisfy $f \boxtimes \zeta = g$ if and only if $f = g \boxtimes \mu$ and this is the case if and only if

$$\varphi_f(z(1+\varphi_g(z))) = \varphi_g(z). \tag{2.7}$$

We will encounter applications of these formulae in section 4.

2.4. **The weak singleton condition.** We can now formulate the problem considered in this paper. De Finetti's theorem states that any exchangeable sequence of classical random variables can be realized as a conditional i.i.d. sequence. There is no general noncommutative analog of this; however since amalgamated freeness is the free analog of conditional independence, one may ask for conditions under which exchangeability systems can be written in the form (2.3). We have already seen that the vanishing of crossing cumulants is a necessary condition. Another necessary condition is the weak singleton condition to be defined now.

Definition 2.5 ([BS96]). (a) An exchangeability system $\mathcal{E} = (\mathcal{U}, \varphi, \mathcal{J})$ satisfies the *singleton condition* if

$$\varphi(X_1^{(i_1)} X_2^{(i_2)} \cdots X_n^{(i_n)}) = 0$$

whenever one of the indices i_j appears only once and the corresponding random variable X_j satisfies $\varphi(X_j) = 0$.

(b) An exchangeability system $\mathcal{E}$ satisfies the *weak singleton condition* if

$$\varphi(X_1^{(i_1)} X_2^{(i_2)} \cdots X_n^{(i_n)}) = 0$$

whenever one of the indices i_j appears only once and the corresponding random variable X_j satisfies $\varphi(X_j^{(1)^*} X_j^{(2)}) = 0$.

The weak singleton condition is indeed weaker than the singleton condition, because it follows from Corollary 2.12 that the condition $\varphi(X^{(1)^*} X^{(2)}) = 0$ implies $\varphi(X) = 0$. The next proposition shows that a weak singleton condition holds also for cumulants, if it holds for moments.

Proposition 2.6. *Let $\mathcal{E}$ be an exchangeability system in which the weak singleton condition holds. Then $K_\pi^{\mathcal{E}}(X_1, X_2, \dots, X_n) = 0$ whenever the partition π contains a singleton $\{j\}$ such that $\varphi(X_j^{(1)^*} X_j^{(2)}) = 0$.*

Proof. Indeed,

$$K_\pi^{\mathcal{E}}(X_1, X_2, \dots, X_n) = \sum_{\sigma \leq \pi} \varphi_\sigma(X_1, X_2, \dots, X_n)\, \mu(\sigma, \pi)$$

and all terms vanish, because each $\sigma \leq \pi$ contains the singleton $\{j\}$. ☐

The starting point of this paper is the following observation and its corollary.

Lemma 2.7. *Let $\pi \in \Pi_n$ be an alternating partition, i.e., a partition in which neighbouring elements are in different blocks. Then any noncrossing partition $\sigma \leq \pi$ contains at least a singleton.*

Proof. Any noncrossing partition σ contains at least one interval block and the condition $\sigma \leq \pi$ implies that this interval block has length 1, i.e., it is a singleton. ☐

Corollary 2.8. *A noncrossing exchangeability system which satisfies the singleton condition is given by a reduced free product.*

Proof. Let $X_1, \dots, X_n \in \mathcal{A}$ with $\varphi(X_j) = 0$ and let $h(1)$, $h(2)$, $\dots$, $h(n)$ be indices such that $h(j) \neq h(j+1)$. We have to show that $\varphi(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = 0$. The singleton condition implies that $\varphi_\pi(X_1, X_2, \dots, X_n) = 0$ whenever π contains a singleton and consequently for any such π the corresponding cumulant $K_\pi^{\mathcal{E}}(X_1, X_2, \dots, X_n)$ vanishes. Now

$$\varphi(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = \sum_{\pi \leq \ker h} K_\pi^{\mathcal{E}}(X_1, X_2, \dots, X_n)$$

and by assumption the sum extends over noncrossing partitions only. By Lemma 2.7 any such partition contains a singleton and the corresponding cumulant vanishes because of Proposition 2.6. □

2.5. **Conditional Expectations.** The proof of 2.8 stays essentially the same if the singleton condition is replaced by the weak singleton condition, resulting in an amalgamated free product. The main problem is the construction of a conditional expectation ψ onto a certain algebra $\mathcal{B}$ and to prove faithfulness of φ on $\mathcal{B}$ in order to apply Lemma 2.7.
The construction of ψ is the same as in the commutative case.

Definition 2.9. Let $\mathcal{E} = (\mathcal{U}, \varphi, J)$ be an exchangeability system for some noncommutative probability space $(\mathcal{A}, \varphi)$. We define the conditional expectations ψ_N, $N \in \mathbf{N}$, by

$$\psi_N(X) = \frac{1}{N!} \sum_{\sigma \in \mathfrak{S}_N} \sigma(X);$$

for polynomials, i.e., elements of the form $X = X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}$ this is

$$\psi_N(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = \frac{1}{N!} \sum_{\sigma \in \mathfrak{S}_N} X_1^{(\sigma(h(1)))} X_2^{(\sigma(h(2)))} \cdots X_n^{(\sigma(h(n)))}$$

Note that for fixed h and N large enough, this only depends on $\ker h$.

We collect a few elementary properties of ψ_N. Proofs are easy and can be found in [AL93].

Proposition 2.10. *(1) $\varphi \circ \psi_N = \varphi$.*
(2) $\psi_N(X) = X$ if and only if $\sigma(X) = X$ $\forall \sigma \in \mathfrak{S}_N$.
(3) $\psi_N \circ \psi_M = \psi_N$ for $M \leq N$.
(4) $\psi_N \circ \iota_k = \frac{1}{N} \sum_{j=1}^N \iota_j$ if $k \leq N$.

In contrast to finite exchangeability systems, there is a nonnegative bilinear form available in the infinite case.

Proposition 2.11. *The sesquilinear form*

$$\langle X, Y \rangle = \varphi(Y^{(1)*} X^{(2)})$$

is nonnegative on $\mathcal{A}$.

Proof. Consider for fixed $N \in \mathbf{N}$ the nonnegative expectation

$$\varphi(\psi_N(X^{(1)})^* \psi_N(X^{(1)})) = \left(\frac{1}{N!}\right)^2 \sum_{\sigma, \sigma'} \varphi(X^{(\sigma(1))*} X^{(\sigma'(1))})$$

$$
\begin{aligned}
&= \frac{1}{N^2} \sum_{i,j=1}^{N} \varphi(X^{(i)^*} X^{(j)}) \\
&= \frac{N(N-1)}{N^2} \varphi(X^{(1)^*} X^{(2)}) + \frac{1}{N} \varphi(X^{(1)^*} X^{(1)}).
\end{aligned}
$$

Now letting $N \to \infty$ yields the claim. □

Positivity implies the Cauchy-Schwarz inequality.

Corollary 2.12. *For any* $X, Y \in \mathcal{A}$

$$
\left|\varphi(Y^{(1)^*} X^{(2)})\right| \leq \varphi(X^{(1)^*} X^{(2)})^{1/2} \, \varphi(Y^{(1)^*} Y^{(2)})^{1/2}
$$

Similarly one can prove a multivariable Cauchy-Schwarz inequality, cf. Remark 2.3.

Corollary 2.13. *Let* $\{g(1), g(2), \ldots, g(m)\}$ *and* $\{h(1), h(2), \ldots, h(n)\}$ *be disjoint sets of indices. Then*

$$
\begin{aligned}
&\left|\varphi(Y_n^{(h(n))^*} \cdots Y_1^{(h(1))^*} X_1^{(g(1))} \cdots X_m^{(g(m))})\right|^2 \\
&\quad \leq \varphi(X_m^{(g'(m))^*} \cdots X_1^{(g'(1))^*} X_1^{(g(1))} \cdots X_m^{(g(m))}) \, \varphi(Y_n^{(h(n))^*} \cdots Y_1^{(h(1))^*} Y_1^{(h'(1))} \cdots Y_n^{(h'(n))})
\end{aligned}
$$

where h' *and* g' *are index functions whose ranges are disjoint from the ranges of* h *and* g.

After these preparations we can consider two situations in which the weak singleton condition holds.

Proposition 2.14. *An exchangeability system satisfies the weak singleton condition in the following two situations.*

(a) φ *is faithful and the exchangeability system comes from an amalgamated free product.*
(b) The state is tracial.

Proof. (a) Let ψ be the conditional expectation with respect to which the algebras are free. Then

$$
\begin{aligned}
0 &= \varphi(X^{(1)^*} X^{(2)}) \\
&= \varphi(\psi(X^{(1)^*} X^{(2)})) \\
&= \varphi(\psi(X^{(1)})^* \psi(X^{(1)}))
\end{aligned}
$$

implies that $\psi(X) = 0$ and if X appears as a singleton in some word, then the expectation of the word vanishes. Indeed, if $X_j = X$ and the index $h(j)$ appears only once in the range of the index function h, then we may condition on $\mathcal{A}_{h(j)}$ and obtain

$$
\varphi(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))} = \varphi(X_1^{(h(1))} X_2^{(h(2))} \cdots \psi(X_j^{(h(j))}) \cdots X_n^{(h(n))}) = 0
$$

(b) Let X_1, X_2,… $X_n \in \mathcal{A}$ and $h : [n] \to \mathbf{N}$ be an index function such that $h(j)$ is a singleton and assume that $\varphi(X_j^{(1)^*} X_j^{(2)}) = 0$. We have to show that $\varphi(X_1^{(h(1))} \cdots X_n^{(h(n))})$ vanishes. By traciality we may assume without loss of generality that $j = n$. Then we may apply the Cauchy-Schwarz inequality of Corollary 2.13 and with any index function h' whose range is disjoint from that of h we obtain

$$
\left|\varphi(X_1^{(h(1))} \cdots X_n^{(h(n))})\right|^2 \leq \varphi(X_1^{(h(1))} \cdots X_{n-1}^{(h(n-1))} X_{n-1}^{(h'(n-1))^*} \cdots X_1^{(h'(1))^*}) \, \varphi(X_n^{(h'(n))^*} X_n^{(h(n))})
$$

$$= 0$$

□

2.6. **Statement of main result.** The first part of Proposition 2.14 shows that the weak singleton condition is a necessary condition for an exchangeability system to come from an amalgamated free product. We can now state the main theorem of this paper.

Theorem 2.15. *If a noncrossing exchangeability system $\mathcal{E} = (\mathcal{U}, \varphi, \mathcal{J})$ with faithful state φ satisfies the weak singleton condition, then it can be embedded into an amalgamated free product such that the interchangeable algebras are free with amalgamation over some subalgebra $\mathcal{B}$.*

Remark 2.16. Other examples where crossing cumulants vanish are Boolean independence [SW97] and more generally conditional free independence [BLS96]. In these examples however Theorem 2.15 does not apply because either the state is not faithful or the weak singleton condition fails: Let $(\mathcal{U}, \varphi, \psi) = \bigstar(\mathcal{A}_i, \varphi_i, \psi_i)$ be the conditionally free exchangeability system for $(\mathcal{A}, \varphi)$, cf. [BLS96] or section I.4.7. We only need the defining property

$$\varphi(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = \varphi(X_1)\,\varphi(X_2) \cdots \varphi(X_n)$$

whenever $\psi(X_j) = 0$ and $h(j) \neq h(j+1)$ for every $1 \leq j \leq n-1$, and the modified pyramidal law

$$\varphi(X_1^{(1)} Y^{(2)} X_2^{(1)}) = \varphi(X_1)\,\varphi(Y)\,\varphi(X_2) + \psi(Y)\,(\varphi(X_1 X_2) - \varphi(X_1)\,\varphi(X_2)),$$

cf. [BLS96] or Lemma I.4.14. Let us investigate the weak singleton condition for some element $X \in \mathcal{A}$. Denoting $\overset{\circ}{X} = X - \psi(X)$ we compute

$$\begin{aligned}
\varphi(X^{(1)^*} X^{(2)}) &= \varphi((\overset{\circ}{X}{}^{(1)^*} + \psi(X^*))\,(\overset{\circ}{X}{}^{(2)} + \psi(X))) \\
&= (\varphi(\overset{\circ}{X}{}^{(1)^*} \overset{\circ}{X}{}^{(2)}) + (\varphi(X^*) - \psi(X^*))\,\psi(X) + \psi(X^*)\,(\varphi(X) - \psi(X)) + |\psi(X)|^2 \\
&= |\varphi(X)|^2
\end{aligned}$$

Now let $X \in \mathcal{A}$ be any element with $\varphi(X) = 0$ but $\psi(X) \neq 0$ (this is possible unless $\varphi = \psi$; in the latter case we have just usual freeness) and find elements Y and Z such that $\varphi(YZ) \neq \varphi(Y)\varphi(Z)$. Then

$$\varphi(Y^{(1)} X^{(2)} Z^{(1)}) = \varphi(Y)\,\varphi(X)\,\varphi(Z) + \psi(X)\,(\varphi(YZ) - \varphi(Y)\,\varphi(Z))$$

does not vanish as it should if the weak singleton condition were true. It follows from Proposition 2.14 that the conditional free product cannot be embedded into an amalgamated free product if the state φ is faithful.

3. A de Finetti Lemma

In this section we prove an asymptotic factorization property of the conditional expectations of Definition 2.9. First we need to review noncrossing partitioned conditional expectations [Spe98].

Definition 3.1 ([Spe98]). Let ψ be a conditional expectation. For a noncrossing partition $\pi \in NC_n$ let $b = \{k, k+1, \dots, l\}$ be an interval block and define recursively

$$\psi[\pi](X_1, X_2, \dots, X_n) = \psi[\pi \setminus b](X_1, X_2, \dots, X_{k-1}, \psi(X_k X_{k+1} \cdots X_l) X_{l+1}, \dots, X_n)$$

These partitioned expectations appear in the calculation of the amalgamated free cumulants.

Proposition 3.2 ([Spe98])**.** *Let* $\mathcal{E} = (\bigstar_{\mathcal{B}}\mathcal{A}_i, \tilde{\psi}, \mathcal{J})$ *be the amalgamated free exchangeability system for a* $\mathcal{B}$*-valued noncommutative probability space* $(\mathcal{A}, \psi)$ *and* h *be an index function. Then for any noncrossing partition* ρ *and any finite sequence* $X_1, X_2, \dots, X_n \in \mathcal{A}$ *we have*

$$\psi_\rho(X_1, X_2, \dots, X_n) = \psi[\rho](X_1, X_2, \dots, X_n).$$

Similarly, for an arbitrary index function h *and any noncrossing partition* ρ *s.t.* $\rho \geq \ker h$ *we have*

$$\tilde{\psi}(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = \tilde{\psi}[\rho](X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))})$$

We will show that the conditional expectations of Definition 2.9 asymptotically have the same property. In the proof the following elementary estimate is needed in two places.

Lemma 3.3. *Let* j, p, N *be positive integers with* $j \leq N$ *and* $p \leq N$, *then*

$$1 - \left(1 - \frac{j}{N}\right)\left(1 - \frac{j}{N-1}\right)\cdots\left(1 - \frac{j}{N-p+1}\right) \leq \frac{pj}{N-p+1}$$

Proof. Denote E_p the quantity in question. Clearly the sequence E_p satisfies $0 \leq E_p \leq 1$, is nondecreasing and therefore $E_p \geq E_1 = \frac{j}{N}$. Moreover it satisfies the recursion

$$\begin{aligned} E_{p+1} &= 1 - \left(1 - \frac{j}{N}\right)\left(1 - \frac{j}{N-1}\right)\cdots\left(1 - \frac{j}{N-p}\right) \\ &= E_p + \frac{j}{N-p}(1 - E_p) \end{aligned}$$

We proceed by induction to show that $C_p = \frac{pj}{N-p+1}$ is an upper bound. Suppose that for E_p we the estimate $E_p \leq C_p$ holds. Then

$$\begin{aligned} E_{p+1} &\leq C_p + \frac{j}{N-p}\left(1 - \frac{j}{N}\right) \\ &= \frac{pj}{N-p+1} + \frac{j}{N-p}\left(1 - \frac{j}{N}\right) \\ &\leq \frac{pj}{N-p} + \frac{j}{N-p} \\ &\leq \frac{(p+1)j}{N-p} \end{aligned}$$

□

Lemma 3.4. *Let* $\pi \in \Pi_n$ *and* $\rho \in NC_n$ *s.t.* $\rho \geq \pi$ *containing* $p = |\pi|$ *and* $r = |\rho|$ *blocks, respectively. Then for* $N \geq p$

$$\left\| \psi_N(X_1^{(\pi(1))} X_2^{(\pi(2))} \cdots X_n^{(\pi(n))}) - \psi_N[\rho](X_1^{(\pi(1))} X_2^{(\pi(2))} \cdots X_n^{(\pi(n))}) \right\| \leq \frac{(2r-1)p^2}{N-p+1} \prod \|X_i\|$$

Proof. Let $\rho = \rho_1 < \rho_2 < \cdots < \rho_r = \hat{1}_n$ be a maximal chain in $[\rho, \hat{1}_n] \cap NC_n$, i.e., ρ_k has exactly $r-k+1$ blocks. It will be convenient to label the blocks b_j of ρ in such a way that

$$\rho_k = \{b_1 \cup b_2 \cup \cdots \cup b_k, b_{k+1}, \dots, b_r\}$$

and correspondingly we label the blocks a_j of π in such a way that $a_1, \dots, a_{j_1} \subseteq b_1$, $a_{j_1+1}, \dots, a_{j_2} \subseteq b_2$, etc., $a_{j_{r-1}+1}, \dots, a_{j_r} \subseteq b_r$. Denote $p_k = j_k - j_{k-1}$ the number of blocks of π which are contained in the kth block b_k of ρ. Let $\tilde{\rho} = \rho/\pi$, i.e., the partition of the block set of π induced by ρ: For i, $j \in \{1, \dots, p\}$ we set $i \sim_{\tilde{\rho}} j$ if a_i and a_j are contained in the same block of ρ. We have to compare the first term

$$\begin{aligned}\psi_N(X_1^{(\pi(1))} X_2^{(\pi(2))} \cdots X_n^{(\pi(n))}) &= \frac{1}{N!} \sum_{\sigma \in \mathfrak{S}_N} X_1^{(\sigma(\pi(1)))} X_2^{(\sigma(\pi(2)))} \cdots X_n^{(\sigma(\pi(n)))} \\ &= \frac{(N-p)!}{N!} \sum_{\substack{h:[p]\to[N] \\ \ker h = \hat{0}_p}} X_1^{(h(\pi(1)))} X_2^{(h(\pi(2)))} \cdots X_n^{(h(\pi(n)))}\end{aligned} \tag{3.1}$$

with the second one

$$\begin{aligned}\psi_N[\rho](X_1^{(\pi(1))} X_2^{(\pi(2))} \cdots X_n^{(\pi(n))}) &= \left(\frac{1}{N!}\right)^r \sum_{\sigma_1, \dots, \sigma_r \in \mathfrak{S}_N} X_1^{(\sigma_{\rho(1)}(\pi(1)))} X_2^{(\sigma_{\rho(2)}(\pi(2)))} \cdots X_n^{(\sigma_{\rho(n)}(\pi(n)))} \\ &= \left(\prod_{k=1}^r \frac{(N-p_k)!}{N!}\right) \sum_{\substack{h:[p]\to[N] \\ \ker h \wedge \tilde{\rho} = \hat{0}_p}} X_1^{(h(\pi(1)))} X_2^{(h(\pi(2)))} \cdots X_n^{(h(\pi(n)))}\end{aligned} \tag{3.2}$$

As in the proof of [AL93, Lemma 2.6] we now split (3.2) as

$$\begin{aligned}&\psi_N[\rho](X_1^{(\pi(1))} X_2^{(\pi(2))} \cdots X_n^{(\pi(n))}) \\ &\quad = \left(\prod_{k=1}^r \frac{(N-p_k)!}{N!}\right) \Biggl(\sum_{k=1}^{r-1} \sum_{\substack{h:[p]\to[N] \\ \ker h \wedge \tilde{\rho}_k = \hat{0}_p \\ \ker h \wedge \tilde{\rho}_{k+1} > \hat{0}_p}} X_1^{(h(\pi(1)))} X_2^{(h(\pi(2)))} \cdots X_n^{(h(\pi(n)))} \\ &\qquad\qquad + \sum_{\substack{h:[p]\to[N] \\ \ker h = \hat{0}_p}} X_1^{(h(\pi(1)))} X_2^{(h(\pi(2)))} \cdots X_n^{(h(\pi(n)))}\Biggr)\end{aligned}$$

Up to a multiplicative constant, the last term is the same as (3.1), and we will show that the constants are asymptotically the same; but first we will bound the $r-1$ terms of the sum. The conditions $\ker h \wedge \tilde{\rho}_k = \hat{0}_p$ and $\ker h \wedge \tilde{\rho}_{k+1} > \hat{0}_p$ mean that $h|_{\{1,\dots,j_k\}}$ is injective, but $h|_{\{1,\dots,j_{k+1}\}}$ is not, i.e., at least one of the indices $h(j_k+1), h(j_k+2), \dots, h(j_{k+1})$ is contained in $\{h(1), \dots, h(j_k)\}$; remember that $h|_{\{j_k+1,\dots,j_{k+1}\}}$ *is* injective. Thus

$$\sum_{\substack{h:[p]\to[N] \\ \ker h \wedge \tilde{\rho}_k = \hat{0}_p \\ \ker h \wedge \tilde{\rho}_{k+1} > \hat{0}_p}} = \sum_{h(1),\dots,h(k) \text{ distinct}} \sum_{\substack{h(j_k+1),\dots,h(j_{k+1}) \text{ distinct} \\ \{h(1),\dots,h(k)\} \cap \{h(j_k+1),\dots,h(j_{k+1})\} \neq \emptyset}} \sum_{h(j_{k+2}),\dots,h(j_r)}$$

There are $N(N-1)\cdots(N-j_k+1)$ different choices for $h(1)$, $h(2)$,…,$h(j_k)$,

$$N(N-1)\cdots(N-p_{k+1}+1) - (N-j_k)(N-j_k-1)\cdots(N-j_k-p_{k+1}+1)$$

choices for $h(j_k+1)$, $h(2)$,…,$h(j_{k+1})$, and

$$\prod_{s=k+2}^{r} \frac{N!}{(N-p_s)!}$$

possibilities to choose the remaining indices $h(j_{k+1}+1)$,…,$h(p)$. The kth term can therefore be estimated by

$$\left(\prod_{s=1}^{k+1} \frac{(N-p_s)!}{N!}\right) \frac{N!}{(N-j_k)!} \left(\frac{N!}{(N-p_{k+1})!} - \frac{(N-j_k)!}{(N-j_k-p_{k+1})!} \right) \prod_{j=1}^{n} \|X_j\|$$

By Lemma 3.3

$$\begin{aligned}
&\frac{(N-p_{k+1})!}{N!} \left(\frac{N!}{(N-p_{k+1})!} - \frac{(N-j_k)!}{(N-j_k-p_{k+1})!} \right) \\
&\qquad = 1 - \frac{(N-j_k)(N-j_k-1)\cdots(N-j_k-p_{k+1}+1)}{N(N-1)\cdots(N-p_{k+1}+1)} \\
&\qquad = 1 - \left(1 - \frac{N-j_k}{N} \frac{N-j_k-1}{N-1} \cdots \frac{N-j_k-p_{k+1}+1}{N-p_{k+1}+1}\right) \\
&\qquad = 1 - \left(1 - \frac{j_k}{N}\right)\left(1 - \frac{j_k}{N-1}\right) \cdots \left(1 - \frac{j_k}{N-p_{k+1}+1}\right) \\
&\qquad \le \frac{p_{k+1} j_k}{N-p_{k+1}+1}
\end{aligned}$$

and therefore the kth term is smaller than

$$\begin{aligned}
&\frac{N(N-1)\cdots(N-j_k+1)}{N(N-1)\cdots(N-p_1+1)\cdots N(N-1)\cdots(N-p_k+1)} \frac{p_{k+1} j_k}{N-p_{k+1}+1} \prod \|X_i\| \\
&\qquad\qquad \le \frac{p_{k+1} j_k}{N-p_{k+1}+1} \prod \|X_i\|
\end{aligned}$$

and

$$\sum_{k=1}^{r-1} \frac{p_{k+1} j_k}{N-p_{k+1}+1} \le (r-1) \frac{\bar{p} p}{N-\bar{p}+1}$$

where $\bar{p} = \max p_k$. Now we come to the difference between the final term and (3.1).

$$\begin{aligned}
&\left\| \left(\prod_{k=1}^{r} \frac{(N-p_k)!}{N!} - \frac{(N-p)!}{N!} \right) \sum_{\ker h = \hat{0}_p} X_1^{(h(\pi(1)))} X_2^{(h(\pi(2)))} \cdots X_n^{(h(\pi(n)))} \right\| \\
&\qquad \le \left(\prod_{k=1}^{r} \frac{(N-p_k)!}{N!} - \frac{(N-p)!}{N!} \right) N(N-1)\cdots(N-p+1) \prod \|X_i\| \\
&\qquad = \left(1 - \prod_{k=1}^{r} \frac{(N-p_k)!}{N!} N(N-1)\cdots(N-p+1) \right) \prod \|X_i\| \\
&\qquad = \left(1 - \prod_{k=1}^{r} \frac{(N-j_{k-1})(N-j_{k-1}-1)\cdots(N-j_k+1)}{N(N-1)\cdots(N-p_k+1)} \right) \prod \|X_i\|
\end{aligned}$$

$$
\begin{aligned}
&= \left(1 - \prod_{k=1}^{r}\left(1 - \frac{j_{k-1}}{N}\right)\left(1 - \frac{j_{k-1}}{N-1}\right)\cdots\left(1 - \frac{j_{k-1}}{N-p_k+1}\right)\right)\prod \|X_i\| \\
&\leq \left(1 - \prod_{k=1}^{r}\left(1 - \frac{p_k j_{k-1}}{N-p_k+1}\right)\right)\prod \|X_i\| \\
&\leq \left(1 - \left(1 - \frac{p\bar{p}}{N-\bar{p}+1}\right)^{r}\right)\prod \|X_i\| \\
&\leq r\frac{p\bar{p}}{N-\bar{p}+1}\prod \|X_i\|
\end{aligned}
$$

by Lemma 3.3. □

The conditional expectations ψ_N need not converge, but we can construct a limit by extending the algebra with the help of the GNS-construction as in [AL93]. Let $\pi : \mathcal{U} \to B(\mathfrak{H})$ be the GNS representation of $\mathcal{U}$ on $\mathfrak{H} = L^2(\mathcal{U}, \varphi)$. By assumption it is faithful and cyclic with cyclic vector ξ_0, i.e., $\{\pi(X)\,\xi_0 : X \in \mathcal{U}\}$ is a dense subspace of $\mathfrak{H}$ and $\varphi(X) = \langle \pi(X)\,\xi_0, \xi_0\rangle$. The action of SG_∞ on $\mathcal{U}$ can be extended to a representation U_σ on $\mathfrak{H}$ which is characterized by

$$U_\sigma \pi(X)\,\xi_0 = \pi(\sigma(X))\,\xi_0 \qquad \forall \sigma \in \mathfrak{S}_\infty \forall X \in \mathcal{U}$$

Let

$$\mathfrak{H}_\infty = \{\xi \in \mathfrak{H} : U_\sigma \xi = \xi \quad \forall \sigma \in \mathfrak{S}_\infty\}$$

be the fixed space of U and $P_\infty : \mathfrak{H} \to \mathfrak{H}_\infty$ the orthogonal projection, and define

$$\psi_\infty(X) = P_\infty \pi(X) P_\infty.$$

Similarly the projection P_N onto

$$[\pi(\psi_N(\mathcal{U}))\,\xi_0] = \{\xi \in \mathfrak{H} : U_\sigma \xi = \xi \quad \forall \sigma \in \mathfrak{S}_N\}$$

is characterized by the property

$$P_N \pi(X)\,\xi = \pi(\psi_N(X))\,\xi \qquad \forall \sigma \in \mathfrak{S}_N \forall X \in \mathcal{U}$$

Clearly every $P_N \geq P_\infty$ and the sequence P_N is monotonically decreasing to its strong limit P_∞. We continue our work in the extended noncommutative probability space $(\tilde{\mathcal{U}}, \tilde{\varphi})$ generated by $\pi(\mathcal{U})$ and P_∞ with state $\tilde{\varphi}(X) = \langle X\xi_0, \xi_0\rangle$. We may also consider it as an operator-valued noncommutative probability space with the conditional expectation

$$\psi_\infty : \tilde{\mathcal{U}} \to \mathcal{B} = P_\infty \tilde{\mathcal{U}} P_\infty$$

and we have $\tilde{\varphi} = \tilde{\varphi} \circ \psi_\infty$. Moreover,

$$\tilde{\varphi}(\psi_\infty(X_1)\psi_\infty(X_2)\cdots\psi_\infty(X_n)) = \lim_{N\to\infty} \varphi(\psi_N(X_1)\psi_N(X_2)\cdots\psi_N(X_n));$$

in particular, $\tilde{\varphi}$ is a trace if φ is a trace. As a corollary to Lemma 3.4 we have the following generalization of [AL93, Lemma 3.1].

Lemma 3.5. *Let $h : [n] \to \mathbf{N}$ be an index function and let ρ be any noncrossing partition such that $\rho \geq \ker h$. Then for any sequence X_1, X_1,…, $X_n \in \mathcal{A}$ we have the factorization*

$$
\begin{aligned}
\varphi(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) &= \lim_{N\to\infty} \langle \psi_N[\rho](X_1^{(h(1))}, X_2^{(h(2))}, \dots, X_n^{(h(n))})\,\xi_0, \xi_0\rangle \\
&= \langle \psi_\infty[\rho](X_1^{(h(1))}, X_2^{(h(2))}, \dots, X_n^{(h(n))})\,\xi_0, \xi_0\rangle
\end{aligned}
$$

First part of the proof of Theorem 2.15. Let us assume for a moment that $\tilde{\varphi}$ is faithful on $\mathcal{B}$. We show that $\tilde{\mathcal{A}}_i = \pi(\mathcal{A}_i)$ are free with amalgamation over $\mathcal{B}$. Let for $X_j \in \mathcal{A}$ and $\psi_\infty(X_j) = 0$ for $1 \leq j \leq n$ and let $h : [n] \to \mathbf{N}$ be an index function with $h(j) \neq h(j+1)$. We have to show that

$$\psi_\infty(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) = 0$$

By the assumed faithfulness it suffices to show that

$$\tilde{\varphi}(\psi_\infty(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))}) Y) = 0 \qquad \forall Y \in \mathcal{B}$$

and it is enough to consider monomials of the form

$$Y = \psi_\infty(Y_1^{(g(1))} Y_2^{(g(2))} \cdots Y_m^{(g(m))})$$

with $Y_j \in \mathcal{A}$ and g an arbitrary index function. Indeed, any element of $\mathcal{B}$ can be approximated by sums of products of elements like above, and for products we have for any $X \in \mathcal{B}$

$$\begin{aligned}\varphi(X \psi_\infty(Y_1^{(f(1))} Y_2^{(f(2))} \cdots Y_p^{(f(p))})\, \psi_\infty(Z_1^{(g(1))} Z_2^{(g(2))} \cdots Z_q^{(g(q))}))&\\ = \varphi(X \psi_\infty(Y_1^{(f(1))} Y_2^{(f(2))} \cdots Y_p^{(f(p))} Z_1^{(g'(1))} Z_2^{(g'(2))} \cdots Z_q^{(g'(q))}))&\end{aligned}$$

where g' is an index function with $\ker g' = \ker g$ and whose range is disjoint from the range of f. Thus consider

$$\begin{aligned}&\tilde{\varphi}(\psi_\infty(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))})\, \psi_\infty(Y_1^{(g(1))} Y_2^{(g(2))} \cdots Y_m^{(g(m))}))\\ &\quad= \lim_{N\to\infty} \tilde{\varphi}(\psi_N(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))})\, \psi_N(Y_1^{(g(1))} Y_2^{(g(2))} \cdots Y_m^{(g(m))}))\\ &\quad= \lim_{N\to\infty} \tilde{\varphi}(\psi_N(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))} Y_1^{(g(1))} Y_2^{(g(2))} \cdots Y_m^{(g(m))}))\end{aligned}$$

where we assume without loss of generality that h and g have disjoint range

$$\begin{aligned}&\quad= \tilde{\varphi}(X_1^{(h(1))} X_2^{(h(2))} \cdots X_n^{(h(n))} Y_1^{(g(1))} Y_2^{(g(2))} \cdots Y_m^{(g(m))})\\ &\quad= \sum_{\substack{\rho_1 \leq \ker h\\ \rho_2 \leq \ker g}} K^{\mathcal{E}}_{\rho_1 \cup \rho_2}(X_1, \dots, X_n, Y_1, \dots, Y_m)\end{aligned}$$

and by assumption the sum runs over all noncrossing partitions only. By lemma 2.7 any noncrossing partition $\rho_1 \leq \ker h$ contains a singleton, say $\{j\}$. Now

$$\begin{aligned}0 &= \tilde{\varphi}(\psi_\infty(X_j)^* \psi_\infty(X_j))\\ &= \lim_{N\to\infty} \tilde{\varphi}(\psi_N(X_j)^* \psi_N(X_j))\\ &= \lim_{N\to\infty} \tilde{\varphi}(\psi_N(X_j^{(1)^*} X_j^{(2)}))\\ &= \tilde{\varphi}(X_j^{(1)^*} X_j^{(2)})\end{aligned}$$

and we may apply Proposition 2.6 to every term of the sum to see that it vanishes. □

The main problem is now to prove faithfulness of $\tilde{\varphi}$ on $\mathcal{B}$. In the tracial case we may dispose of this problem as follows.

End of the proof of Theorem 2.15 in the tracial case. If φ is a trace, so is $\tilde{\varphi}$ and its kernel is a twosided ideal. Since φ is faithful, the intersection of $\pi(\mathcal{U})$ with $\ker \tilde{\varphi}$ is trivial and therefore $\mathcal{U}$ is faithfully embedded into the quotient algebra $\tilde{U}/\ker \tilde{\varphi}$, on which the trace is faithful. □

In the non-tracial case there is more work to do, namely we will show that the extended state $\tilde{\varphi}$ is indeed faithful on $\mathcal{B}$. This is done by showing that in the case of noncrossing cumulants a very strong law of large numbers holds.

4. A NONCOMMUTATIVE L^p-INEQUALITY IN THE CASE OF NONCROSSING CUMULANTS

Our aim is to show that $\psi_\infty(X) = 0$ if $\varphi(X^{(1)^*}X^{(2)}) = 0$. That is, such random variables satisfy a very strong law of large numbers. We need a combinatorial proof in order to use the combinatorial information about cumulants, that is, we will use the fact that for a faithful state we have

$$\left\|\sum X_i\right\| = \lim_{p\to\infty} \varphi(((\sum X_i)^*(\sum X_i))^p))^{1/2p} \tag{4.1}$$

The proof is somewhat in the spirit of [Pis00] where it is shown that the noncommutative L^p-norms of so-called p-orthogonal sums (of which our situation is a special case) can be estimated

$$\left\|\sum X_i\right\|_{L^{2p}(\tau)} \leq \frac{3\pi}{2}\, p\, S(X,p)$$

where

$$S(X,p) = \max\left\{\left\|\left(\sum X_i^* X_i\right)^{1/2}, \left(\sum X_i X_i^*\right)^{1/2}\right\|\right\}$$

However in order to get something useful out of (4.1) we will need constants which stay bounded as p tends to infinity. This is related to the question in [Pis00, Remark 0.3] whether there are uniform constants for free martingale inequalities, owing to the fact that the size of the lattice NC_n of noncrossing partitions is of order 4^n, while the size of the lattice of all partitions Π_n is much bigger. The tracial version in Proposition 6.1 gives further evidence for a positive answer to this question. For our purposes however we need a variant of the inequality for i.i.d. sequences also in the nontracial case.

Proposition 4.1. *Assume that a noncrossing exchangeability system $\mathcal{E}$ satiesfies the weak singleton condition and has a faithful state. Then for any selfajoint random variable X with $\varphi(X^{(1)}X^{(2)}) = 0$ the interchangeable sequence $X^{(i)}$ satisfies the inequality*

$$\left\|\sum_{i=1}^N X^{(i)}\right\| \leq \frac{2\sqrt{N}}{1-\frac{1}{\sqrt{N}}}\,\|X\|$$

for every $N \geq 2$.

Proof. We give three estimates with increasing difficulty and accuracy. Roughly the idea is as follows. We assume that $X^{(i)}$ are as in the statement of the proposition. By faithfulness of φ, we can use (4.1) although the "L^p-norm" associated to φ is not really a norm. We can expand the latter in terms of cumulants:

$$\begin{aligned}\varphi(\left(\sum X^{(i)}\right)^p) &= \sum_{\pi\in NC_p} K_\pi(\sum X^{(i)}) \\ &= \sum_{\pi\in NC_p}\sum_{\ker h\geq\pi} K_\pi(X^{(h(1))}, X^{(h(2))}, \dots, X^{(h(p))}) \\ &= \sum_\pi N^{|\pi|} K_\pi(X)\end{aligned}$$

First estimate. Because of the weak singleton condition only partitions without singletons contribute. Any such partition has at most $\frac{p}{2}$ blocks and therefore the sum is of order $N^{p/2}$ times the number of noncrossing partitions:

$$\leq N^{p/2}\frac{1}{p+1}\binom{2p}{p}\max_{\pi}\lvert K_\pi(X)\rvert$$

Each cumulant K_π in turn can be estimated by

$$\begin{aligned}\lvert K_\pi(X)\rvert &= \left\lvert\sum_{\sigma\leq\pi}\varphi_\sigma(X)\,\mu_{NC}(\sigma,\pi)\right\rvert\\ &\leq \frac{1}{p+1}\binom{2p}{p}\lVert X\rVert^p\max_{\sigma,\pi}\lvert\mu_{NC}(\sigma,\pi)\rvert\\ &\simeq 16^p\lVert X\rVert^p\end{aligned}$$

Thus by this first rough estimate we obtain the inequality

$$\left\lvert\varphi\left((\sum X^{(i)})^p\right)\right\rvert\leq 64^pN^{p/2}\lVert X\rVert^p$$

and taking limits

$$\left\lVert\sum X^{(i)}\right\rVert\leq 64\sqrt{N}\lVert X\rVert\,.$$

Second estimate. With a little effort, we can improve on the constant considerably. First note that we can evaluate $a_\pi=\sum_{\sigma\leq\pi}\lvert\mu_{NC}(\sigma,\pi)\rvert$ explicitly. Since μ_{NC} is a multiplicative function, so are $\lvert\mu_{NC}\rvert$ and $a=\lvert\mu_{NC}\rvert\boxtimes\zeta$. By applying the Kreweras complementation map we have

$$a_n=\sum_{\sigma\in NC_n}\left\lvert\mu_{NC}(\sigma,\hat{1}_n)\right\rvert=\sum_{\sigma\in NC_n}\left\lvert\mu_{NC}(\hat{0}_n,\sigma)\right\rvert \tag{4.2}$$

i.e., $a=\lvert\mu_{NC}\rvert\boxtimes\zeta$ and we can use (2.7). The characteristic series of $\lvert\mu_{NC}\rvert$ is

$$\varphi_{\lvert\mu\rvert}=\frac{1}{2}(1-\sqrt{1-4z})$$

and $\varphi_a(z)$ satisfies the equation

$$\frac{1}{2}\left(1-\sqrt{1-4z(1+\varphi_a(z))}\right)=\varphi_a(z).$$

Together with the condition $\varphi_a(0)=0$ this yields the solution

$$\varphi_a(z)=\frac{1}{2}(1-z-\sqrt{1-6z+z^2})=z+2z^2+6z^3+22z^4+\dots.$$

This is the generating function of the "large Schröder numbers" [Sta99, Deu01].
We have to estimate

$$\left\lvert\sum_{\pi\in NC_p^{\geq2}}N^{\lvert\pi\rvert}K_\pi(X)\right\rvert\leq\sum_{\pi\in NC_p^{\geq2}}N^{\lvert\pi\rvert}a_\pi\lVert X\rVert^p$$

where $NC_p^{\geq2}$ is the set of noncrossing partitions without singletons. The sequence

$$b_n=\sum_{\pi\in NC_n^{\geq2}}N^{\lvert\pi\rvert}a_\pi$$

is the characteristic sequence of the convolution of the multiplicative function $N \cdot \overset{\circ}{a}$ with characteristic sequence $(N\overset{\circ}{a}_n)_n$ with the ζ-function, where

$$\overset{\circ}{a}_n = \begin{cases} 0 & n = 1 \\ a_n & n \geq 2 \end{cases}$$

and

$$\varphi_{\overset{\circ}{a}}(z) = \frac{1}{2}\left(1 - 3z - \sqrt{1 - 6z + z^2}\right)$$

The characteristic series $\varphi_b(z)$ can be found by yet another appeal to (2.7), namely it satisfies the equation

$$N\varphi_{\overset{\circ}{a}}(z(1 + \varphi_b(z))) = \varphi_b(z)$$

and the relevant solution is

$$\varphi_b(z) = \frac{2(N+1)}{N + 2 + 3Nz + N\sqrt{1 - 6z + (1 - 8N)z^2}} - 1.$$

The dominant singularity comes from the radicand $1 - 6z + (1 - 8N)z^2$. The zeros of the latter are $\frac{1}{8N-1}\left(\pm\sqrt{2(N+1)} - 3\right)$ and therefore

$$b_n \sim \left(\frac{8N - 1}{2\sqrt{2(N+1)} - 3}\right)^n;$$

it follows that

$$\left\|\sum X^{(i)}\right\| \leq \frac{8N - 1}{2\sqrt{2(N+1)} - 3}\,\|X\|\,.$$

The constant tends to $2\sqrt{2}$ as $N \to \infty$, which is not bad, as the best possible constant is 2.

Third estimate. With even some more effort, one can obtain the optimal constant (at least as $N \to \infty$) as follows. The previous estimate was done using the numbers a_n from (4.2) and we neglected the fact that for the calculation of the cumulants $K_\pi(X)$ partitions with singletons do not contribute. Thus it will be more accurate to work with the numbers

$$\tilde{a}_n = \sum_{\pi \in NC_n^{\geq 2}} \left|\mu_{NC}(\pi, \hat{1}_n)\right| \tag{4.3}$$

which constitute the characteristic sequence of the multiplicative function $\overset{\circ}{\zeta} \mathbin{\boxed{\star}} |\mu_{NC}|$ where

$$\overset{\circ}{\zeta}_n = \begin{cases} 0 & n = 1 \\ 1 & n \geq 2 \end{cases}$$

is the Zeta function on the poset of noncrossing partitions without singletons. This convolution can be carried out with the aid of (2.6). The “Fourier transforms” of the functions

$$\varphi_{\overset{\circ}{\zeta}}(z) = \sum_{n=2}^{\infty} z^n = \frac{z^2}{1 - z} \qquad \text{and} \qquad \varphi_{|\mu|}(z) = \frac{1}{2}\left(1 - \sqrt{1 - 4z}\right)$$

are

$$\mathcal{F}_{\overset{\circ}{\zeta}} = \frac{\pm\sqrt{z^2 + 4z} - z}{2z} \qquad \text{and} \qquad \mathcal{F}_{|\mu|}(z) = 1 - z$$

respectively. Therefore

$$\mathcal{F}_{\overset{\circ}{\zeta}\boxed{\star}|\mu|}(z) = \frac{\pm\sqrt{z^2+4z}-z}{2z}\,(1-z)$$

i.e., $y = y(z) = \varphi_{\overset{\circ}{\zeta}\boxed{\star}|\mu|}(z)$ satisfies the algebraic equation

$$y(1-y)(1-y-z) = z^2.$$

We are interested in the asymptotics of the numbers

$$\tilde{b}_n = \sum_{\pi\in NC_n} N^{|\pi|}\tilde{a}_\pi$$

whose generating function can be determined by (2.7), namely

$$N\varphi_{\tilde{a}}(z(1+\varphi_{\tilde{b}}(z))) = \varphi_{\tilde{b}}(z)$$

Thus $x = x(z) = \varphi_{\tilde{b}}(z)$ satisfies the equations

$$N\varphi_{\tilde{a}}(z(1+x)) = x \qquad \frac{x}{N}(1-\frac{x}{N})(1-\frac{x}{N}-z(1+x)) = z^2(1+x)^2$$

therefore $x = x(z)$ is the solution of the equation

$$g(x,z) = \frac{x}{N}\,\left(1-\frac{x}{N}\right)\,\left(1-\frac{x}{N}-z\,(x+1)\right) - z^2\,(x+1)^2 = 0 \tag{4.4}$$

If $x(z)$ has a singularity at z, then both $g(x,z)=0$ and $\partial_x g(x,z)=0$ and therefore z is a zero of the resultant

$$Res(g(x,z),\partial_z g(x,z)) = \frac{1}{N^9}(N\,z+1)\,(N+1)^2\,z^2$$
$$\times\left(8\,z-2\,N\,z+32\,z^2-4\,z^3+26\,N\,z^2+16\,N\,z^3-N^2\,z^2+10\,N^2\,z^3-N^2\,z^4+4\,N^3\,z^4-5\right),$$

cf. [CLO98, FS01]. By Pringsheim's theorem we know that the dominant singularity is positive and therefore it must be a root of the last factor

$$r(z) = -5+(8-2N)\,z+(32+26N-N^2)\,z^2+(-4+16N+10N^2)\,z^3+(-N^2+4N^3)\,z^4$$

We claim that $r(z)\neq 0$ for $0\leq z\leq \frac{1}{2\sqrt{N}}\left(1-\frac{1}{\sqrt{N}}\right)$. Indeed, let $z = \frac{\alpha}{2\sqrt{N}}\left(1-\frac{1}{\sqrt{N}}\right)$ with $0\leq\alpha\leq 1$, then it is tedious but not difficult to verify that

$$\begin{aligned}
&r(\frac{\alpha}{2\sqrt{N}}\left(1-\frac{1}{\sqrt{N}}\right)) \\
&\qquad = N\left(\frac{\alpha^4}{4}-\frac{\alpha^2}{4}\right)+N^{1/2}\left(-\alpha+\frac{\alpha^2}{2}+\frac{5}{4}\,\alpha^3-\alpha^4\right) \\
&\qquad\quad -5+\alpha+\frac{25}{4}\,\alpha^2-\frac{15}{4}\,\alpha^3+\frac{23}{16}\,\alpha^4+N^{-1/2}(4\alpha-13\alpha^2+\frac{23}{4}\,\alpha^3-\frac{3}{4}\,\alpha^4) \\
&\qquad\quad +N^{-1}(-4\alpha+\frac{29}{2}\,\alpha^2-\frac{29}{4}\,\alpha^3-\frac{1}{8}\,\alpha^4)+N^{-3/2}(-16\alpha^2+\frac{11}{2}\,\alpha^3+\frac{1}{8}\,\alpha^4) \\
&\qquad\quad +N^{-2}(8\,\alpha^2-\frac{1}{2}\,\alpha^3-\frac{1}{16}\,\alpha^4)-N^{-5/2}\frac{3}{2}\,\alpha^3+N^{-3}\frac{1}{2}\,\alpha^3 \\
&\qquad = -\frac{N}{4}\,\alpha^2(1-\alpha^2)-N^{1/2}\left(\frac{1}{4}(\frac{1}{4}-(\alpha-\frac{1}{2})^2)+(1-\alpha)+\alpha(\alpha-\frac{1}{2})^2\right)
\end{aligned}$$

$$
\begin{aligned}
&- (1 - N^{-1/2}) \left(\frac{1}{16} + (1-\alpha)\left(\frac{47}{16} + \frac{29}{8}\alpha + 2(1-\alpha)^2 + \frac{23}{16}(1-(1-\alpha)^3)\right)\right) \\
&- (N^{-1/2} - N^{-1})\Biggl(\frac{877}{256} + \frac{3}{32}(1-\alpha) + \frac{27}{8}\left(\frac{1}{8} - \left(\alpha - \frac{1}{2}\right)^3\right) \\
&\qquad\qquad + \left(\alpha-\frac{1}{2}\right)^2\left(\frac{685}{256} + \frac{11}{16}\left(\frac{1}{4} - \left(\alpha-\frac{1}{2}\right)^2\right)\right)\Biggr) \\
&- N^{-1}\left(\frac{15}{16} + 3(1-\alpha)(1-(1-\alpha)^2) + \frac{37}{8}(1-\alpha)^2 - \frac{9}{16}(1-\alpha)^4\right) \\
&- (N^{-3/2} - N^{-2})\alpha^2(16 - \frac{11}{2}\alpha - \frac{1}{4}\alpha^2) - N^{-2}\alpha^2(8 - 5\alpha - \frac{3}{16}\alpha^2) \\
&- N^{-5/2}(3 - N^{-1/2})\frac{1}{2}\alpha^3
\end{aligned}
$$

which is strictly negative for $0 \le \alpha \le 1$. Therefore asymptotically as n tends to infinity we have

$$\tilde{b}_n \le \left(\frac{2\sqrt{N}}{1 - \frac{1}{\sqrt{N}}}\right)^n .$$

□

End of the proof of Theorem 2.15 in the nontracial case. It remains to prove faithfulness of the state $\tilde{\varphi}$ on $\mathcal{B}$. Let $X \in \mathcal{B}$ such that $\tilde{\varphi}(X^*X) = 0$, say $X = \psi_\infty(W)$, where $W \in \mathcal{U}$. By remark 2.3 it is enough to consider $W = Y^{(1)}$ with $Y \in \mathcal{A}$. By assumption

$$
\begin{aligned}
0 &= \tilde{\varphi}(\psi_\infty(Y^{(1)})^*\psi_\infty(Y^{(1)})) \\
&= \tilde{\varphi}(\psi_\infty({Y^{(1)}}^* Y^{(2)})) \\
&= \varphi({Y^{(1)}}^* Y^{(2)})
\end{aligned}
$$

and by Proposition 4.1 this implies that $\psi_\infty(Y^{(1)}) = 0$. □

5. Weak freeness

In this section we discuss the notion of weak freeness, which together with the weak singleton condition implies vanishing of crossing cumulants.

Definition 5.1. Let $\mathcal{E} = (\mathcal{U}, \tilde{\varphi}, \mathcal{J})$ be an exchangeability system for a noncommutative probability space $(\mathcal{A}, \varphi)$. For an index set $I \subseteq \mathbf{N}$ denote $\mathcal{A}_I$ the subalgebra of $\mathcal{U}$ generated by $(\mathcal{A}_i)_{i\in I}$. We say that $\mathcal{E}$ satisfies *weak freeness* if

$$\varphi(X_1 X_2 \cdots X_n) = 0$$

whenever $\varphi({X_j^{(1)}}^* X_j^{(2)}) = 0$, I_k are disjoint index sets and $X_j \in \mathcal{A}_{(I_{i_j})}$ with $i_j \ne i_{j+1}$ for every j. Here $X_j^{(1)}$ and $X_j^{(2)}$ refer to copies of X_j in $A_{I'_{i_j}}$ and $A_{I''_{i_j}}$, where I'_{i_j} and I''_{i_j} are disjoint copies of I_{i_j}.

It will be convenient to adapt the exchangeability system as follows. Decompose $\mathbf{N}$ into an infinite union of disjoint copies of itself $\mathbf{N} = \bigcup_{j=0}^{\infty} I_j$. Then relabel the indices and consider the exchangeability system with embeddings $\iota_{ij} : \mathcal{A} \to \mathcal{A}_{i,j} \subseteq \mathcal{U}$, $i, j \in \mathbf{N}$. Thus $\mathcal{U}$ is also an exchangeability system for $\tilde{A} = \bigvee_{j\in\mathbf{N}} \mathcal{A}_{0j}$ and we will work with this interpretation in this section, i.e., our random variables X are elements of $\tilde{\mathcal{A}}$ and $X^{(i)}$ are elements of $\tilde{\mathcal{A}}_i = \bigvee_{j\in\mathbf{N}} \mathcal{A}_{ij}$. Thus if $X = X_1^{(0,j_1)} X_2^{(0,j_2)} \cdots X_n^{(0,j_n)}$, then $X^{(i)} = X_1^{(i,j_1)} X_2^{(i,j_2)} \cdots X_n^{(i,j_n)}$. The weak freeness condition of Definition 5.1 can be rephrased more clearly as follows, namely

$$\tilde{\varphi}(X_1^{(i_1)} X_2^{(i_2)} \cdots X_n^{(i_n)}) = 0$$

whenever $X_j \in \tilde{\mathcal{A}}$ with $\tilde{\varphi}(X_j^{(1)^*} X_j^{(2)}) = 0$ and $i_j \neq i_{j+1}$ for all $1 \leq j \leq n-1$.
We need this regrouping in order to define an asymptotic conditional expectation which is used to transfer proofs from the amalgamated free situation. As in section 2.5, we define symmetrizing maps

$$\begin{aligned} \psi_N : \tilde{\mathcal{A}} &\to \tilde{\mathcal{A}} \\ X &\mapsto \frac{1}{N!} \sum_{\sigma\in\mathfrak{S}_N} \sigma(X) \end{aligned}$$

which will allow us to construct asymptotically ψ-centered random variables.

Lemma 5.2. *Let $X_j \in \tilde{\mathcal{A}}$ be polynomials, that is, linear combinations of elements of the form*

$$Z_1^{(0,j_1)} Z_2^{(0,j_2)} \cdots Z_m^{(0,j_m)}$$

with $Z_j \in \mathcal{A}$. Then

$$K_\pi(X_1, \ldots, X_{k-1}, \psi_N(X_k), X_{k+1}, \ldots, X_n) \xrightarrow[N\to\infty]{} 0$$

unless $\{k\}$ is a singleton of π. In other words, $\psi_N(X_k)$ is asymptotically independent from the rest.

Proof. Indeed

$$K_\pi(X_1, \ldots, X_{k-1}, \psi_N(X_k), X_{k+1}, \ldots, X_n) = \frac{1}{N!} \sum_{\sigma\in\mathfrak{S}_N} K_\pi(X_1, \ldots, X_{k-1}, \sigma(X_k), X_{k+1}, \ldots, X_n)$$

Let s be the maximal superscript appearing in the polynomials X_j. If σ maps each $1 \leq j \leq s$ to some index strictly greater than s, then $\sigma(X_k)$ is independent from the other X_j and by assumption the cumulant vanishes. The number of permutations σ of this type is $(N-s)(N-s-1)\cdots(N-2s)\cdot(N-s)!$, i.e., almost all permutations, because the ratio of the rest is

$$\frac{N! - (N-s)(N-s-1)\cdots(N-2s)\cdot(N-s)!}{N!} \xrightarrow[N\to\infty]{} 0.$$

□

When calculating cumulants of elements $X_j \in \tilde{\mathcal{A}}$, we can thus replace X_j by $X_j - \psi_N(X_j)$ for each non-singleton index j and then let N tend to infinity. Replacing X_j by $X_j - \psi_N(X_j)$ allows to apply the weak singleton condition, because due to the permutation invariance of the state we have

$$\tilde{\varphi}(\sigma(X)^{(1)^*} X^{(2)}) = \tilde{\varphi}(X^{(1)^*} X^{(2)})$$

$$\begin{aligned}
\tilde\varphi((X^{(1)} - \psi_N(X)^{(1)})^*(X^{(2)} - \psi_N(X)^{(2)}))& \\
&= \tilde\varphi(X^{(1)*}(X)^{(2)}) - \frac{1}{N!}\sum_\sigma \tilde\varphi(\sigma(X)^{(1)*}X^{(2)}) \\
&\quad - \frac{1}{N!}\sum_\sigma \tilde\varphi(X^{(1)*}\sigma(X)^{(2)}) + \frac{1}{(N!)^2}\sum_{\sigma,\tau} \tilde\varphi(\sigma(X)^{(1)*}\tau(X)^{(2)}) \\
&= 0
\end{aligned}$$

Theorem 5.3. *Let $\mathcal{E} = (\mathcal{U}, \tilde\varphi, \mathcal{J})$ be an exchangeability system for a noncommutative probability space $(\mathcal{A}, \varphi)$ with faithful state $\tilde\varphi$ such that both weak freeness and the weak singleton condition holds. Then crossing cumulants vanish. In particular, $\mathcal{E}$ can be embedded into an amalgamated free product.*

Proof. The proof consists of three parts by reducing an arbitrary crossing partition to an alternating partition without singletons.
We will use the following terminology. A partition $\pi \in \Pi_n$ is called *alternating* if $i \not\sim_\pi i+1$ for all $1 \le i \le n-1$, that is, adjacent elements are in different blocks of π. For non-alternating partitions we denote by

$$\operatorname{cn}(\pi) = \#\{(k, k+1) : k \sim_\pi k+1, 1 \le k \le n-1\}$$

the *number of connected neighbours* of π. Clearly π is alternating if and only if $\operatorname{cn}(\pi) = 0$.
Step 1. Alternating partitions without singletons. First assume that π is an alternating partition without singletons. Let $\varepsilon > 0$. Then by Lemma 5.2 we may find $N > 0$ such that

$$K_\pi(X_1, \dots, X_n) = K_\pi(X_1 - \psi_N(X_1), \dots, X_n - \psi_N(X_n)) + R_n$$

with error term $|R_n| < \varepsilon$. Now a look at the moment-cumulant formula

$$K_\pi(X_1 - \psi_N(X_1), \dots, X_n - \psi_N(X_n)) = \sum_{\sigma \le \pi} \varphi_\sigma(X_1 - \psi_N(X_1), \dots, X_n - \psi_N(X_n))\, \mu(\sigma, \pi)$$

shows that the sum runs over alternating partitions and by weak freeness every term vanishes.
Step 2. Reducing everything to alternating partitions. The aim is now to express an arbitrary cumulant in terms of alternating ones. We will use the product formula of Leonov and Shiryaev from Proposition 2.2 to reduce the number of connected neighbours. Consider a partition $\pi \in \Pi_n$ with crossings and $\operatorname{cn}(\pi) > 0$. Pick an arbitrary element k with $k \sim_\pi k+1$. We will express K_π as a sum of cumulants K_ρ s.t. $k \not\sim_\rho k+1$ and $\operatorname{cn}(\rho) < \operatorname{cn}(\pi)$. Let $\hat\pi = \pi/[k = k+1] \in \Pi_{n-1}$ be the partition obtained from π by identifying k and $k+1$. Then by Proposition 2.2 we have with $\nu = \{\{1\}, \{2\}, \dots, \{k, k+1\}, \dots, \{n\} = \mid \mid \dots \mid \sqcap \mid \dots \mid$ the decomposition

$$\begin{aligned}
K_{\hat\pi}(X_1, \dots, X_k X_{k+1}, \dots, X_n) &= \sum_{\rho \vee \nu = \pi} K_\rho(X_1, \dots, X_k, X_{k+1}, \dots, X_n) \\
&= K_\pi(X_1, \dots, X_n) + \sum_{\substack{\rho \vee \nu = \pi \\ \rho < \pi}} K_\rho(X_1, \dots, X_n)
\end{aligned}$$

Each contributing partition ρ in the second sum is obtained from π by splitting the block containing k and $k+1$ into two in such a way that k and $k+1$ are separated. In particular,

$\operatorname{cn}(\rho) \leq \operatorname{cn}(\pi) - 1$ and $\operatorname{cn}(\hat{\pi}) = \operatorname{cn}(\pi) - 1 < \operatorname{cn}(\pi)$ as well. To conclude, we have

$$K_\pi(X_1, \dots, X_n) = K_{\hat{\pi}}(X_1, \dots, X_k X_{k+1}, \dots, X_n) - \sum_{\substack{\rho \vee \nu = \pi \\ \rho < \pi}} K_\rho(X_1, \dots, X_n)$$

and all partitions appearing on the right hand side have less connected neighbours than π. Repeating this operation finitely many times we end up with a linear combination of alternating partitions (possibly involving singletons).

Step 3. Getting rid of singletons. Assume that after step 2 we have arrived at an alternating partition π. Then for N large enough, we have by Lemma 5.2 again

$$K_\pi(X_1, \dots, X_n) \approx K_\pi(\tilde{X}_1, \dots, \tilde{X}_n)$$

where

$$\tilde{X}_j = \begin{cases} X_j & \text{if } \{j\} \text{ is a singleton of } \pi \\ X_j - \psi_N(X_j) & \text{if } \{j\} \text{ is not a singleton of } \pi \end{cases}$$

We may therefore assume without loss of generality that $\psi(X_j^{(1)^*} X_j^{(2)}) = 0$ for the non-singleton indices j. If the singleton entries k satisfy $\psi(X_k^{(1)^*} X_k^{(2)}) = 0$ as well, then we may proceed as in step one. If however there are singletons for which this is not the case, we may eliminate them as follows. Let $\{k\}$ be the first of these critical singletons, then we may write the cumulant as

$$K_\pi(X_1, \dots, X_n) = K_\pi(X_1, \dots, X_k - \psi_N(X_k), \dots, X_n) + K_\pi(X_1, \dots, \psi_N(X_k), \dots, X_n)$$

The first term has one critical singleton less than the left hand side and the second term can be treated with the product formula as follows. Let again $\hat{\pi} = \pi/[k = k+1] \in \Pi_{n-1}$ be the partition obtained from π by identifying k with $k+1$. Moreover put $\nu = \{\{1\}, \{2\}, \dots, \{k, k+1\}, \dots, \{n\}\} = \mid\ \mid\ \dots \mid \sqcap \mid\ \dots \mid$ and let $\tilde{\pi} = \pi \vee \nu$ be the partition obtained from π by adjoining the singleton $\{k\}$ to the block containing $k+1$. Then we have by Proposition 2.2 again

$$\begin{aligned} K_{\hat{\pi}}(X_1, X_2, \dots, \psi_N(X_k) X_{k+1}, \dots, X_n) \\ &= K_{\tilde{\pi}}(X_1, X_2, \dots, \psi_N(X_k), X_{k+1}, \dots, X_n) \\ &\quad + \sum_{\substack{\rho \vee \nu = \tilde{\pi} \\ \rho < \tilde{\pi}}} K_\rho(X_1, X_2, \dots, \psi_N(X_k), X_{k+1}, \dots, X_n) \end{aligned}$$

$K_{\tilde{\pi}}$ vanishes asymptotically by Lemma 5.2 and so do all K_ρ in which $k \sim_\pi k+1$, and hence the only nontrivial term on the right hand side is the cumulant indexed by $\rho = \pi$, because this is the only one in which k is a singleton. Thus

$$K_\pi(X_1, X_2, \dots, \psi_N(X_k), X_{k+1}, \dots, X_n) \approx K_{\hat{\pi}}(X_1, X_2, \dots, \psi_N(X_k) X_{k+1}, \dots, X_n)$$

and $\hat{\pi}$ has one singleton less than π. Repeating this procedure we end up with a linear combination of alternating partitions without singletons and step one of the proof applies. □

6. Appendix: Free L^p Inequalities

We consider now the tracial version of Proposition 4.1. In this case Hölder's inequality is available and we can get estimates in terms of the L^p-norms.

Proposition 6.1. *Assume that an noncrossing exchangeability system $\mathcal{E} = (\mathcal{U}, \tau, \mathcal{J})$ is tracial and faithful. Then for any sequence of $\mathcal{E}$-independent random variables X_i with $\tau(X_i^{(1)^*} X_i^{(2)}) = 0$ the inequality*

$$\left\| \sum_{i=1}^{N} X_i \right\|_{L^{2p}(\tau)} \leq C_{2p} S(X, 2p)$$

holds with $C_{2p} \leq \frac{3\pi}{4z_0} \simeq 9.85859$ as $p \to \infty$ where z_0 is computed in (6.1).

Proof. We expand the L^p-norm in terms of cumulants:

$$\tau\left(\left(\sum X_i^* X_j\right)^p\right) = \sum_{\pi \in NC_{2p}} K_\pi(\sum X_i^*, \sum X_i, \ldots, \sum X_i^*, \sum X_i)$$

and for each π, we estimate the cumulant K_π. Because of the weak singleton condition, only partitions without singletons are involved in the sums.

$$\begin{aligned} &K_\pi(\sum X_i^*, \sum X_i, \ldots, \sum X_i^*, \sum X_i) \\ &\quad= \sum_{\ker h \geq \pi} K_\pi(X_{h(1)}^*, X_{h(2)}, \ldots, X_{h(2p-1)}^*, X_{h(2p)}) \\ &\quad= \sum_{\ker h \geq \pi} \sum_{\sigma \leq \pi} \tau_\sigma(X_{h(1)}^*, X_{h(2)}, \ldots, X_{h(2p-1)}^*, X_{h(2p)})\, \mu_{NC}(\sigma, \pi) \end{aligned}$$

Now for fixed σ we have [Pis00, Sublemma 3.3 and Lemma 3.4]

$$\left| \sum_{\ker h \geq \pi} \tau_\sigma(X_{h(1)} X_{h(2)} \cdots X_{h(2p)}) \right| \leq \left\| \sum \lambda(g_i) \otimes X_i \right\|_{2p}^{2p} \leq \left(\frac{3\pi}{4}\right)^{2p} S(X, 2p)$$

where $\lambda(g_i)$ is the left regular representation of the generators of the free group. Indeed, we can find a suitable discrete group G and elements $F_1, \ldots, F_{2p}$ in $L^p(\tau_G \otimes \tau)$ such that

$$\|F_k\|_{2p} = \left\| \sum \lambda(g_i) \otimes X_i \right\|_{2p}$$

and

$$\sum_{\ker h \geq \pi} \tau_\sigma(X_{h(1)} X_{h(2)} \cdots X_{h(2p)}) = \tau_G \otimes \varphi(F_1 F_2 \cdots F_{2p}).$$

The construction is done as in the proof of [Pis00, Sublemma 3.3], with the slight modification that X_i is replaced by $X_i^{(\sigma(k))}$ when constructing F_k. With these preparations we continue similarly as before

$$\tau\left(\left(\sum X_i^* X_j\right)^p\right) \leq \sum_{\pi \in NC^{\geq 2}} \sum_{\substack{\sigma \in NC_{2p}^{\geq 2} \\ \sigma \leq \pi}} |\mu_{NC}(\sigma, \pi)| \left(\frac{3\pi}{4}\right)^{2p} S(X, 2p)$$

We know the asymptotics of

$$\tilde{b}_n = \sum_{\pi\in NC^{\geq 2}} \sum_{\substack{\sigma\in NC_{2p}^{\geq 2}\\ \sigma\leq\pi}} |\mu_{NC}(\sigma,\pi)|$$

from its generating function, which satisfies the equation (4.4) with $N = 1$. Its dominant singularity is a zero of the resultant

$$r(z) = 3z^4 + 22z^3 + 57z^2 + 6z - 5$$

and the singularity in question is

$$z_0 = -\frac{11}{6} + \frac{1}{6}\sqrt{7+\gamma} + \sqrt{14 - \gamma + \frac{992}{\sqrt{7+\gamma}}} \simeq 0.238999 \tag{6.1}$$

where

$$\gamma = 9\left(207 - 48\sqrt{3}\right)^{\frac{1}{3}} + 9\left(207 + 48\sqrt{3}\right)^{\frac{1}{3}};$$

consequently $\tilde{b}_n \simeq z_0^{-n}$ as $n \to \infty$. □

7. A Free Analog of Brillinger's Formula

7.1. **Brillinger's formula for classical cumulants.** Let $(\Omega, \mathcal{A}, \mu)$ be a probability space with expectation $\mathbf{E}\,X = \int f\,d\mu$. Classical independence corresponds to the exchangeability system $L^\infty(\Omega^\infty, \mu^\infty)$ for the algebra $L^\infty(\Omega, \mu)$ of bounded random variables. For a finite sequence of random variables $X_1, X_2, \ldots, X_n$ on Ω the partitioned moment functional (2.2) completely factors

$$\mathbf{E}_\pi(X_1, X_2, \ldots, X_n) = \prod_j \mathbf{E} \prod_{i\in\pi_j} X_i$$

and similarly the partitioned cumulants factor as

$$\kappa_\pi(X_1, X_2, \ldots, X_n) = \sum_{\substack{\sigma\in\Pi_n\\ \sigma\leq\pi}} \mathbf{E}_\sigma(X_1, X_2, \ldots, X_n)\,\mu(\pi, \hat{1}_n) = \prod_{B\in\pi} \kappa_{|B|}(X_i : i \in B)$$

where μ is the Möbius function on the partition lattice Π_n.

Similarly taking conditional expectations with respect to a σ-subalgebra $\mathcal{B} \subseteq \mathcal{A}$, the partitioned conditional expectations are the $\mathcal{B}$-measurable random variables

$$\mathbf{E}_\pi(X_1, X_2, \ldots, X_n|\mathcal{B}) = \prod_j \mathbf{E}(\prod_{i\in\pi_j} X_i|\mathcal{B})$$

and the *conditioned cumulants* are the $\mathcal{B}$-measurable random variables

$$\kappa_\pi(X_1, X_2, \ldots, X_n|\mathcal{B}) = \sum_{\sigma\leq\pi} \mathbf{E}_\sigma(X_1, X_2, \ldots, X_n|\mathcal{B})\,\mu(\pi, \hat{1}_n)$$

The conditioned cumulants are again multiplicative on blocks and can be used to detect conditional independence, namely if $X_1, X_2, \ldots, X_n$ can be divided into two groups which are mutually independent conditionally on $\mathcal{B}$, then the cumulant $\kappa_n(X_1, X_2, \ldots, X_n)$ vanishes.

The starting point of this paper is the following formula, due to Brillinger [Bri69]:

$$\kappa_n(X_1, X_2, \ldots, X_n) = \sum_{\pi\in\Pi_n} \kappa_{|\pi|}(\kappa_{|\pi_j|}(X_i : i \in \pi_j|\mathcal{B}) : j = 1, \ldots, |\pi|) \tag{7.1}$$

where for a partition $\pi = \{B_1, B_2, \dots, B_p\} \in \Pi_n$ we denote by $|\pi| = p$ its *size*.
The aim of this paper is to establish an analogue of this formula for free cumulants by adapting a lattice theoretical proof of the above formula which is due to Speed [Spe83].

7.2. **Free cumulants.**

Definition 7.1 ([Voi95])**.** Let $(\mathcal{A}, \varphi)$ be a noncommutative $\mathcal{B}$-valued probability space; i.e., $\mathcal{A}$ is a unital complex algebra, $\mathcal{B} \subseteq \mathcal{A}$ is a unital subalgebra and $\varphi : \mathcal{A} \to \mathcal{B}$ is a conditional expectation. Subalgebras $\mathcal{A}_i$ which contain $\mathcal{B}$ are called *free with amalgamation over* $\mathcal{B}$ if

$$\varphi(X_1 X_2 \cdots X_n) = 0$$

whenever $X_j \in \mathcal{A}_{i_j}$, $\varphi(X_j) = 0$ and $i_j \neq i_{j+1}$ $\forall j$. When $\mathcal{B} = \mathbf{C}$, we recover the definition of *freeness*.

7.3. **Conditioned free cumulants.** Suppose we are given algebras $\mathcal{C} \subseteq \mathcal{B} \subseteq \mathcal{A}$ and conditional expectations $\mathcal{A} \xrightarrow{\psi} \mathcal{B} \xrightarrow{\varphi} \mathcal{C}$. We identify φ with $\varphi \circ \psi : \mathcal{A} \to \mathcal{C}$ and wish to express the $\mathcal{C}$-valued cumulants C^φ in terms of the $\mathcal{B}$-valued cumulants C^ψ.

Definition 7.2. We define a partitioned moment function φ of the partitioned cumulants C_π^ψ, namely for $\sigma \geq \pi$ we define $\varphi_\sigma \circ C_\pi^\psi(X_1, X_2, \dots, X_n)$ recursively as follows. Let $\sigma_j = \{k+1, \dots, l\}$ be an interval block of σ and $\pi|_{\sigma_j} = \{\pi_{i_1}, \pi_{i_2}, \dots, \pi_{i_m}\}$ the blocks of π which are contained in σ_j, then we put

$$\begin{aligned}&\varphi_\sigma \circ \psi_\pi(X_1, X_2, \dots, X_n)\\ &\qquad = \varphi_{\sigma \setminus \{\sigma_j\}} \circ \psi_{\pi \setminus \pi|_{\sigma_j}}(X_1, X_2, \dots, X_k, \varphi(\psi_{\pi|_{\sigma_j}}(X_{k+1}, \dots, X_l)) X_{l+1}, X_{l+2}, \dots, X_n)\end{aligned}$$

and

$$\varphi_\sigma \circ C_\pi^\psi(X_1, X_2, \dots, X_n) = \sum_{\tau \leq \pi} \varphi_\sigma \circ \psi_\tau(X_1, X_2, \dots, X_n)\, \mu(\tau, \pi).$$

By multiplicativity we have

$$\begin{aligned}&\varphi_\sigma \circ C_\pi^\psi(X_1, X_2, \dots, X_n)\\ &\qquad = \varphi_{\sigma \setminus \{\sigma_j\}} \circ C^\psi_{\pi \setminus \pi|_{\sigma_j}}(X_1, X_2, \dots, X_k, \varphi(C^\psi_{\pi|_{\sigma_j}}(X_{k+1}, \dots, X_l)) X_{l+1}, X_{l+2}, \dots, X_n).\end{aligned}$$

and by Möbius inversion and the invariance $\varphi = \varphi \circ \psi$ we have

$$\varphi_\pi(X_1, X_2, \dots, X_n) = \sum_{\sigma \leq \pi} \varphi_\pi \circ C_\sigma^\psi(X_1, X_2, \dots, X_n).$$

Now we apply the cumulant construction in each block of σ to define "cumulants of cumulants":

$$C_\sigma^\varphi \circ C_\pi^\psi(X_1, X_2, \dots, X_n) = \sum_{\pi \leq \rho \leq \sigma} \varphi_\rho \circ C_\pi^\psi(X_1, X_2, \dots, X_n)\, \mu(\rho, \sigma)\,.$$

In total this means that

$$C_\sigma^\varphi \circ C_\pi^\psi(X_1, X_2, \dots, X_n) = \sum_{\tau \leq \pi} \sum_{\pi \leq \rho \leq \sigma} \varphi_\rho \circ \psi_t(X_1, X_2, \dots, X_n)\, \mu(\rho, \sigma)\, \mu(\tau, \pi)$$

This function is multiplicative on the blocks and we have by Möbius inversion

$$\varphi_\sigma \circ C_\pi^\psi(X_1, X_2, \dots, X_n) = \sum_{\pi \leq \rho \leq \sigma} C_\rho^\varphi \circ C_\pi^\psi(X_1, X_2, \dots, X_n)$$

Here is now the analogue of (7.1) for free cumulants. As expected, noncrossing partitions appear, but we also have to take care of noncommutativity.

Theorem 7.3.

$$C_n^{\varphi}(X_1, X_2, \dots, X_n) = \sum_{\sigma \in NC_n} C_n^{\varphi} \circ C_\sigma^{\psi}(X_1, X_2, \dots, X_n)$$

Proof. The proof of [Spe83] can be repeated literally after replacing the lattice Π_n by its sublattice NC_n:

$$\begin{aligned} C_n^{\varphi}(X_1, X_2, \dots, X_n) &= \sum_{\pi \in NC_n} \varphi_\pi(X_1, X_2, \dots, X_n)\, \mu(\pi, \hat{1}_n) \\ &= \sum_{\pi \in NC_n} \sum_{\sigma \leq \pi} \sum_{\sigma \leq \rho \leq \pi} C_\rho^{\varphi} \circ C_\sigma^{\psi}(X_1, X_2, \dots, X_n)\, \mu(\pi, \hat{1}_n) \\ &= \sum_{\pi \in NC_n} \sum_{\rho \in NC_n} \sum_{\sigma \in NC_n} C_\rho^{\varphi} \circ C_\sigma^{\psi}(X_1, X_2, \dots, X_n)\, \zeta(\sigma, \rho)\, \zeta(\rho, \pi)\, \mu(\pi, \hat{1}_n) \\ &= \sum_{\rho \in NC_n} \sum_{\sigma \leq \rho} C_\rho^{\varphi} \circ C_\sigma^{\psi}(X_1, X_2, \dots, X_n)\, \delta(\rho, \hat{1}_n) \end{aligned}$$

□

7.4. **An application.** As an application we reprove a characterization of freeness from the recent paper [NSS02]. To illustrate our approach, let us first give a proof of a more or less trivial formula from the latter paper.

Proposition 7.4 ([NSS02, Theorem 3.1])**.** *If the ψ-valued cumulants of $X_1, X_2, \dots, X_n$ satisfy*

$$C_k^{\psi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k}) \in \mathcal{C}$$

for all choices of indices $i_1, i_2, \dots, i_k$ and elements $c_1, \dots, c_{k-1} \in \mathcal{C}$, then actually

$$C_k^{\psi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k}) = C_k^{\varphi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k})$$

Proof. By Theorem 7.3 we can expand the φ-cumulant in terms of the ψ-cumulants

$$C_n^{\varphi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k}) = \sum_{\pi} C_{|\pi|}^{\varphi} \circ C_\pi^{\psi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k})$$

where by definition

$$\begin{aligned} &C_{|\pi|}^{\varphi} \circ C_\pi^{\psi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k}) \\ &\qquad = \sum_{\sigma \geq \pi} \varphi_\sigma \circ C_\pi^{\psi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k})\, \mu(\sigma, \hat{1}_n). \end{aligned}$$

By assumption,

$$\varphi_\sigma \circ C_\pi^{\psi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k}) = C_\pi^{\psi}(X_{i_1} c_1, X_{i_2} c_2, \dots, X_{i_{k-1}} c_{k-1}, X_{i_k})$$

and $\sum_{\sigma \geq \pi} \mu(\sigma, \hat{1}_n) = 0$ unless $\pi = \hat{1}_n$. Therefore only the summand corresponding to $\pi = \hat{1}_n$ is nonzero. □

Theorem 7.5 ([NSS02, Theorem 3.6])**.** *Let $\mathcal{C} \subseteq \mathcal{N} \subseteq \mathcal{A}$ be a subalgebra and assume that $\varphi : \mathcal{B} \to \mathcal{C}$ is faithful. Then $\mathcal{N}$ is free from $\mathcal{B}$ over $\mathcal{C}$ if and only if for all finite sequences $X_i \in \mathcal{N}$ and for all $b_i \in \mathcal{B}$ the equality*

$$C_n^\psi(X_1b_1, X_2b_2, \dots, X_{n-1}b_{n-1}, X_n) = \varphi(C_n^\psi(X_1\varphi(b_1), X_2\varphi(b_2), \dots, X_{n-1}\varphi(b_{n-1}), X_n))$$

holds. This is equivalent to saying that for all finite sequences $X_i \in \mathcal{N}$ and for all $b_i \in \mathcal{B}$ we have

$$C_n^\psi(X_1b_1, X_2b_2, \dots, X_{n-1}b_{n-1}, X_n) = C_n^\varphi(X_1\varphi(b_1), X_2\varphi(b_2), \dots, X_{n-1}\varphi(b_{n-1}), X_n).$$

Proof. Assume that the factorization formula holds. Let $X_1, X_2, \dots, X_n \in \mathcal{N}$, $b_1, \dots, b_n \in \mathcal{B}$ s.t. $\varphi(X_i) = 0$ and $\varphi(b_i) = 0$ (or $b_0 = 1$ or $b_1 = 1$ is also allowed). We must show that $\varphi(b_0X_1b_1X_2 \cdots X_nb_n) = 0$. To this end we expand the expectation into ψ-cumulants

$$\varphi(b_0X_1b_1 \cdots X_nb_n)\varphi(\varphi(b_0X_1b_1 \cdots X_nb_n)) = \sum_{\pi \in NC_n} \varphi(C_\pi^\psi(b_0X_1b_1, X_2b_2, \dots, X_nb_n))$$

and $C_\pi^\psi(b_0X_1b_1, X_2b_2, \dots, X_nb_n)) = 0$ for each π because each π has a block which is an interval say of length m starting at some k and the corresponding cumulant contributes the factor

$$C_m^\psi(X_kb_k, X_{k+1}b_{k+1}, \dots X_l) = \varphi(C_m^\psi(X_k\varphi(b_k), X_{k+1}\varphi(b_{k+1}), \dots X_l))$$

which vanishes: if $m \geq 2$ then there is a factor $\varphi(b_k) = 0$ and if $m = 1$, then the term is simply $C_1^\psi(X_k) = \varphi(C_1^\psi(X_k)) = \varphi(X_k) = 0$. Note that we did not need faithfulness of φ for this implication.

For the converse we can use the same argument as in [NSS02], in which a reference algebra $\mathcal{N}'$ is constructed which is also free from $\mathcal{B}$ over $\mathcal{C}$ and which satisfies the cumulant factorization condition and has the same distribution as $\mathcal{N}$. It then follows that $\mathcal{N}$ satisfies the cumulant factorization condition as well.

If you feel like this is cheating, here is a sketch of a direct proof using conditioned cumulants. By faithfulness it suffices to prove that for all finite sequences of random variables $X_i \in \mathcal{N}$ and $b_i \in \mathcal{B}$ we have that

$$\begin{aligned}\varphi(C_n^\psi(X_1b_1, X_2b_2, \dots, X_{n-1}b_{n-1}, X_nb_n))\\ = \varphi(C_n^\psi(X_1\varphi(b_1), X_2\varphi(b_2), \dots, X_{n-1}\varphi(b_{n-1}), X_n\varphi(b_n)))\end{aligned}$$

and that all this is then equal to

$$C_n^\varphi(X_1\varphi(b_1), X_2\varphi(b_2), \dots, X_{n-1}\varphi(b_{n-1}), X_n\varphi(b_n)).$$

In the rest of this proof, we use the notation $\pi\tilde{\cup}\sigma$ for the *interweaved union* of two partitions π and σ of the same order, i.e., the partition of doubled order obtained by alternatingly arranging the points of π and σ on a line. If π and σ are noncrossing and $\sigma \leq K(\pi)$, then the resulting partition is again noncrossing. Indeed by definition $K(\pi)$ is the maximal partition σ s.t. $\pi\tilde{\cup}\sigma$ is noncrossing.

We proceed by induction and compare the formulae

$$\begin{aligned}&C_n^\varphi(X_1b_1, X_2b_2, \dots, X_nb_n)\\ &\quad= \sum_{\pi \in NC_n} C^\varphi_{\pi\tilde{\cup}K(\pi)}(X_1, b_1, X_2, b_2, \dots, X_n, b_n)\end{aligned}$$

$$= \underbrace{C^{\varphi}_{\hat{1}_n \breve{\cup} \hat{0}_n}(X_1, b_1, X_2, b_2, \ldots, X_n, b_n)}_{= C^{\varphi}_n(X_1\varphi(b_1), \ldots, X_n\varphi(b_n))} + \sum_{\pi < \hat{1}_n} C^{\varphi}_{\pi \breve{\cup} K(\pi)}(X_1, b_1, X_2, b_2, \ldots, X_n, b_n)$$

and

$$\begin{aligned} &C^{\varphi}_n(X_1b_1, X_2b_2, \ldots, X_nb_n) \\ &\quad = \sum_{\pi \in NC_n} C^{\varphi}_n \circ C^{\psi}_{\pi}(X_1b_1, X_2b_2, \ldots, X_nb_n) \\ &\quad = \varphi(C^{\psi}_n(X_1b_1, X_2b_2, \ldots, X_nb_n)) + \sum_{\pi < \hat{1}_n} C^{\varphi}_n \circ C^{\psi}_{\pi}(X_1b_1, X_2b_2, \ldots, X_nb_n) \end{aligned}$$

We will prove inductively for $\pi < \hat{1}_n$ the identity

$$C^{\varphi}_n \circ C^{\psi}_{\pi}(X_1b_1, X_2b_2, \ldots, X_nb_n) = C^{\varphi}_{\pi \breve{\cup} K(\pi)}(X_1, b_1, X_2, b_2, \ldots, X_n, b_n). \tag{7.2}$$

Note that the intervals $[\,\pi, \hat{1}_n\,]$ and $[\,0, K(\pi)\,]$ are antiisomorphic via the Kreweras complement. On the other hand, each interval $[\,\pi, \sigma\,]$ is isomorphic to a direct product of full noncrossing partition lattices of various orders, which themselves are antiisomorphic to themselves. Therefore $[\,\pi, \sigma\,]$ is antiisomorphic to itself and finally we get that $[\,\pi, \hat{1}_n\,]$ and $[\,0, K(\pi)\,]$ are also isomorphic.
Here is an explicit isomorphism. Draw π and all the points of $K(\pi)$ between the points of π. Every $\sigma \geq \pi$ is obtained from π by connecting some of its blocks. To this connections correspond connections of $K(\pi)$ as follows. There are two possible relative positions of two blocks of π:

(a) 

(b) 

In both cases connecting the two blocks of π corresponds to connecting the points marked with "×" in the complement.
To prove (7.2), we compare

$$C^{\varphi}_n \circ C^{\psi}_{\pi}(X_1b_1, X_2b_2, \ldots, X_nb_n) = \sum_{\rho \geq \pi} \varphi_{\rho} \circ C^{\psi}_{\pi}(X_1b_1, X_2b_2, \ldots, X_nb_n)\mu(\rho, \hat{1}_n)$$

with

$$\begin{aligned} &C_{\pi \breve{\cup} K(\pi)}(X_1, b_1, X_2, b_2, \ldots, X_n, b_n) \\ &\qquad = \sum_{\sigma_2 \leq K(\pi)} \sum_{\sigma_1 \leq \pi} \varphi_{\sigma_1 \breve{\cup} \sigma_2}(X_1, b_1, X_2, b_2, \ldots, X_n, b_n)\, \mu(\sigma_1, \pi)\, \mu(\sigma_2, K(\pi)). \end{aligned}$$

By induction hypothesis, when $\pi < \hat{1}_n$, we can replace certain elements b_i by their expectation $\varphi(b_i)$ and C^{ψ}_k by C^{φ}_k and we have

$$\sum_{\sigma_1 \leq \pi} \varphi_{\sigma_1 \breve{\cup} \sigma_2}(X_1, b_1, X_2, b_2, \ldots, X_n, b_n)\, \mu(\sigma_1, \pi) = \varphi_{\rho} \circ C^{\psi}_{\pi}(X_1b_1, X_2b_2, \ldots, X_nb_n)$$

where σ_2 is the partition in $[\,\hat{0}_n, K(\pi)\,]$ corresponding to ρ under the isomorphism above. □

Acknowledgements. We are grateful to Roland Speicher for a simplification in the first part of the proof of Theorem 7.5.

References

[AL93] Accardi, L., and Lu, Y. G., *A continuous version of de Finetti's theorem*, Ann. Probab. **21** (1993), 1478–1493.

[BLS96] Bożejko, M., Leinert, M., and Speicher, R., *Convolution and limit theorems for conditionally free random variables*, Pacific J. Math. **175** (1996), 357–388.

[Bri69] Brillinger, D., *The calculation of cumulants via conditioning*, Ann. Inst. Statist. Math. **21** (1969), 375–390.

[BS96] Bożejko, M., and Speicher, R., *Interpolations between bosonic and fermionic relations given by generalized Brownian motions*, Math. Z. **222** (1996), 135–159.

[CLO98] Cox, D., Little, J., and O'Shea, D., *Using algebraic geometry*, Graduate Texts in Mathematics, vol. 185, Springer-Verlag, New York, 1998.

[CT78] Chow, Y. S., and Teicher, H., *Probability theory*, Springer-Verlag, New York, 1978.

[Deu01] Deutsch, E., *A bijective proof of the equation linking the Schröder numbers, large and small*, Discrete Math. **241** (2001), 235–240.

[FS01] Flajolet, P., and Sedgewick, R., *Analytic combinatorics: functional equations, rational, and algebraic functions*, preprint, `http://pauillac.inria.fr/algo/flajolet/Publications/FlSe01.pdf`, 2001, Chapter 8 of forthcoming book "Analytic Combinatorics".

[HM76] Hudson, R. L., and Moody, G. R., *Locally normal symmetric states and an analogue of de Finetti's theorem*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **33** (1975/76), 343–351.

[Hud81] Hudson, R. L., *Analogs of de Finetti's theorem and interpretative problems of quantum mechanics*, Found. Phys. **11** (1981), 805–808.

[Kin78] Kingman, J. F. C., *Uses of exchangeability*, Ann. Probability **6** (1978), 183–197.

[Kre72] Kreweras, G., *Sur les partitions non croisées d'un cycle*, Discrete Math. **1** (1972), 333–350.

[Leh04] Lehner, F., *Cumulants in noncommutative probability theory I. Noncommutative exchangeability systems*, Math. Zeitschr. **248** (2004), no. 1, 67–100, arXiv:math.CO/0210442.

[NS97] Nica, A., and Speicher, R., *A "Fourier transform" for multiplicative functions on non-crossing partitions*, J. Algebraic Combin. **6** (1997), 141–160.

[NSS02] Nica, A., Shlyakhtenko, D., and Speicher, R., *Operator-valued distributions. I. Characterizations of freeness*, Int. Math. Res. Not. (2002), no. 29, 1509–1538, arXiv:math.OA/0201001.

[Pet90] Petz, D., *A de Finetti-type theorem with m-dependent states*, Probab. Theory Related Fields **85** (1990), 65–72.

[Pis00] Pisier, G., *An inequality for p-orthogonal sums in non-commutative L_p*, Illinois J. Math. **44** (2000), 901–923.

[Spe83] Speed, T. P., *Cumulants and partition lattices*, Austral. J. Statist. **25** (1983), 378–388.

[Spe94] Speicher, R., *Multiplicative functions on the lattice of noncrossing partitions and free convolution*, Math. Ann. **298** (1994), 611–628.

[Spe98] Speicher, R., *Combinatorial theory of the free product with amalgamation and operator-valued free probability theory*, Mem. Amer. Math. Soc. **132** (1998), no. 627, x+88.

[Sta86] Stanley, R. P., *Enumerative combinatorics. Vol. I*, The Wadsworth & Brooks/Cole Mathematics Series, Wadsworth & Brooks/Cole Advanced Books & Software, Monterey, CA, 1986, With a foreword by Gian-Carlo Rota.

[Sta99] Stanley, R. P., *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999.

[Stø69] Størmer, E., *Symmetric states of infinite tensor products of C^*-algebras*, J. Functional Analysis **3** (1969), 48–68.

[SW97] Speicher, R., and Woroudi, R., *Boolean convolution*, Free probability theory (Waterloo, ON, 1995), Amer. Math. Soc., Providence, RI, 1997, pp. 267–279.

[Voi95] Voiculescu, D., *Operations on certain non-commutative operator-valued random variables*, Astérisque (1995), no. 232, 243–275, Recent advances in operator algebras (Orléans, 1992).

Franz Lehner, Institut für Mathematik C, Technische Universität Graz, Steyrergasse 30, A-8010 Graz, Austria
E-mail address: `lehner@finanz.math.tu-graz.ac.at`